



OPEN ACCESS

# Clinical characteristics, treatment and outcomes of paediatric COVID-19: a systematic review and meta-analysis

Omar Irfan,<sup>1</sup> Fiona Muttalib,<sup>1</sup> Kun Tang,<sup>1,2</sup> Li Jiang,<sup>1</sup> Zohra S Lassi,<sup>3</sup> Zulfiqar Bhutta <sup>1,4</sup>

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/archdischild-2020-321385>).

<sup>1</sup>Centre for Global Child Health, The Hospital for Sick Children, Toronto, Ontario, Canada

<sup>2</sup>Vanke School of Public Health, Tsinghua University, Beijing, China

<sup>3</sup>Robinson Research Institute, Adelaide Medical School, The University of Adelaide, Adelaide, South Australia, Australia

<sup>4</sup>Institute for Global Health & Development, Aga Khan University, Karachi, Pakistan

## Correspondence to

Dr Zulfiqar Bhutta, Centre for Global Child Health, The Hospital for Sick Children, Toronto, ON M5G 1X8, Canada; [zulfiqar.bhutta@sickkids.ca](mailto:zulfiqar.bhutta@sickkids.ca)

Received 11 December 2020  
Revised 25 January 2021  
Accepted 27 January 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Irfan O, Muttalib F, Tang K, et al. *Arch Dis Child* Epub ahead of print: [please include Day Month Year]. doi:10.1136/archdischild-2020-321385

## ABSTRACT

**Objective** Compare paediatric COVID-19 disease characteristics, management and outcomes according to World Bank country income level and disease severity.

**Design** Systematic review and meta-analysis.

**Setting** Between 1 December 2019 and 8 January 2021, 3350 articles were identified. Two reviewers conducted study screening, data abstraction and quality assessment independently and in duplicate. Observational studies describing laboratory-confirmed paediatric (0–19 years old) COVID-19 were considered for inclusion.

**Main outcomes and measures** The pooled proportions of clinical findings, treatment and outcomes were compared according to World Bank country income level and reported disease severity.

**Results** 129 studies were included from 31 countries comprising 10251 children of which 57.4% were hospitalised. Mean age was 7.0 years (SD 3.6), and 27.1% had a comorbidity. Fever (63.3%) and cough (33.7%) were common. Of 3670 cases, 44.1% had radiographic abnormalities. The majority of cases recovered (88.9%); however, 96 hospitalised children died. Compared with high-income countries, in low-income and middle-income countries, a lower proportion of cases were admitted to intensive care units (ICUs) (9.9% vs 26.0%) yet pooled proportion of deaths among hospitalised children was higher (relative risk 2.14, 95% CI 1.43 to 3.20). Children with severe disease received antimicrobials, inotropes and anti-inflammatory agents more frequently than those with non-severe disease. Subgroup analyses showed that a higher proportion of children with multisystem inflammatory syndrome (MIS-C) were admitted to ICU (47.1% vs 22.9%) and a higher proportion of hospitalised children with MIS-C died (4.8% vs 3.6%) compared with the overall sample.

**Conclusion** Paediatric COVID-19 has a favourable prognosis. Further severe disease characterisation in children is needed globally.

## INTRODUCTION

The coronavirus disease (COVID-19) pandemic caused by severe acute respiratory syndrome coronavirus-2 has spread from a local outbreak in China to a global pandemic within months. On 31 December 2019, a cluster of cases with pneumonia of unknown cause emerged from Wuhan, China. On 30 January 2020, the WHO declared the coronavirus outbreak a Public Health Emergency of

International Concern, and on 11 March 2020, a pandemic. As of 21 January 2021, there have been over 95.6 million confirmed COVID-19 cases and over 2.0 million associated deaths from 216 countries, areas or territories.<sup>1</sup> Children under-19 years of age comprise a small proportion (1%–10%) of the total reported cases<sup>2–5</sup> with a lower risk of developing critical illness from COVID-19 infection compared with adults.<sup>6</sup> Prior systematic reviews of paediatric COVID-19 have described a mild disease in children with good outcomes.<sup>4 7 8</sup> Since the publication of these reviews, the pandemic has spread extensively around the globe. In addition to pulmonary manifestations of COVID-19 in children, reports from Europe, North America, Latin America and Asia have emerged, describing a multi-system inflammatory syndrome children (MIS-C) related to COVID-19 infection.<sup>9–12</sup> COVID-19 has also disrupted essential maternal and child health interventions, including outpatient visits and vaccinations for young children in most countries, further worsening the existing burden on health-care provision and delivery.<sup>13</sup>

The objective of this review, in addition to providing a comprehensive update of the evolving paediatric COVID-19 literature, is a unique comparison of reported cases in low-income and middle-income countries (LMICs) to high-income countries (HICs) and of children with severe versus non-severe disease. Furthermore, the review provides a subgroup analysis of children presenting with symptoms of MIS-C and neonatal cases.

## METHODS

The protocol of the review is registered with PROSPERO (CRD42020183134). This systematic review is reported in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-analysis (PRISMA).

## SEARCH METHODS

The review includes a comprehensive search of MEDLINE, Embase, WHO COVID-19 Database, Chinese COVID-19 Databases (CNKI and Wangfang), Latin-American and Caribbean Health Sciences Literature (LILACS) from 1 December 2019 to 8 January 2021. Complementary searches were conducted in Google Scholar, John Hopkins Health Resource, WHO news and the Chinese and US CDC Library. MedRxiv, BioRxiv and ChinaXiv

were searched for preprints. No language restrictions were applied.

A search strategy was formulated and administered as shown in online supplemental table 1.

### STUDY SELECTION

Observational studies reporting children (0–19 years old) with laboratory-confirmed COVID-19 (serology or RT-PCR) were considered for inclusion. Studies with a subset of children 0–19 years were included if disaggregated data for children were provided. Studies were screened for any overlap in paediatric cases by reviewing institution details and the period reported. Review articles, case reports, commentaries and letters not presenting any original data were excluded. Case reports were excluded to reduce risk of selection bias and over-representation of extreme cases. Covidence Software (2016) was used for screening by two reviewers independently and in duplicate. Key reference lists were screened for additional studies.

### DATA EXTRACTION

Two reviewers conducted data extraction using a prepiloted data form. Data extracted included authors' names, date of publication, study-design, city, country, number of cases, gender, comorbidities, travel and contact history, diagnostic tests for COVID-19, clinical details, laboratory tests, radiological findings, management and outcomes. Disaggregated data by age groups (0–5 years, 5–10 years and >10 years old) and reported disease severity was extracted where available. Criteria for severe disease were as defined within each individual study and included admission to intensive care units (ICUs), use of mechanical ventilation, multiorgan failure and presence of hypoxia (oxygen saturation <92%).

### QUALITY ASSESSMENT

Individual study quality was evaluated independently by the review authors using quality assessment tools developed by the National Heart Lung and Brain Institute (NHLBI)<sup>14</sup> (<https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools>). Study quality was scored out of 8, based on clarity of study objectives, case definition, consecutive inclusion of cases, comparability of included patients, definition and measurement of outcomes, length of follow-up, statistical methods and results. Studies with score 6–8 were considered to be good quality, 4–5 considered fair quality and <4 considered poor quality.

### DATA SYNTHESIS

Categorical data were summarised as counts and proportions. The pooled proportions of reported findings were calculated using Comprehensive Meta-Analysis 2.2.027 using random-effects model.  $I^2$  was calculated to examine statistical heterogeneity ( $I^2 > 50\%$  considered high heterogeneity). The clinical features and outcomes were compared according to (1) World Bank country income level (HICs versus LMICs)<sup>15</sup> and (2) reported disease severity (severe versus non-severe) using pooled proportions and their 95% CIs, supplemented by relative-risk (RR). Subgroup analyses of children with MIS-C and neonatal cases were conducted.

### RESULTS

After removal of duplicates, 3350 citations were screened for inclusion. Full texts of 198 studies were screened and 129 studies<sup>2 3 9–12 16–138</sup> were included (online supplemental figure 1). Sixty-nine studies were excluded as they either presented

overlapping data, did not provide age-disaggregated data for children or were commentaries, editorials or reviews. In terms of study setting, 13 studies were population-based national surveillance studies, 94 studies included only children admitted to hospital and 22 studies reported patients presenting to outpatient clinics or emergency departments (hospitalisation rate of 24.2%, 385/1590).

Sixty studies were from HICs (n=6528) and 69 studies from LMICs (n=3723). Almost one-third of included studies were from China (36/129, 28.0%),<sup>216–3335 37–43 45 52 53 55–57 71 74 138</sup> one-fifth were from the USA (24/129, 18.6%)<sup>3 9 10 47 58 60 64 68–70 79 81 82 85–87 93 98 109 110 115 120 133 136</sup> together comprising almost half of the included sample size (n=4758, 46.4%). The country of origin of included studies and study characteristics are summarised in online supplemental figure 2, tables 2 and 3.

### DEMOGRAPHICS AND EPIDEMIOLOGY

A total of 9335 children from the 129 case series were included in the meta-analysis. Of 8455 children for whom initial disposition was reported, 4851 were hospitalised (57.4%). Among them, 55.5% were men. The patient's ages ranged from 0 to 17 years with mean age of  $7.0 \pm 3.6$  years. Ninety-one of the 129 studies reported age-disaggregated data for infection incidence as shown in online supplemental table 4. Nearly half of the cases were >10 years of age. Contact exposure to COVID-19 was reported in 64.0% of cases. Travel history to an epicentre was reported in 13.0% cases. At-least one underlying comorbidity was reported in 27.1% of cases. The most common reported comorbidities were immunosuppression (15.8%) and lung disease (12.5%).

### CLINICAL MANIFESTATIONS

Table 1 summarises the clinical manifestations reported in the studies. There were 13.1% asymptomatic cases (95% CI 10.4% to 16.3%) who presented primarily through contact exposure in family-clusters (parents, siblings and other relatives). The most common presenting symptoms were fever (63.3%, 95% CI 58.6% to 68.4%) and cough (33.7%, 95% CI 29.6% to 38.1%) followed by nausea or vomiting (20.0%, 95% CI 16.5% to 24.0%) and diarrhoea (19.6%, 95% CI 16.1% to 23.7%). Other symptoms included dyspnoea, nasal-symptoms, rashes, kawasaki-like symptoms, conjunctivitis, fatigue, abdominal pain and neurological symptoms. Sixty-seven of the 129 studies reported age disaggregated data for clinical features (online supplemental table 5). Clinical features were similar in the three age groups:  $\leq 5$  years,  $>5$  to  $\leq 10$  years,  $>10$  years with higher prevalence of abdominal symptoms in children  $>5$  years.

### RADIOLOGICAL AND LABORATORY FINDINGS

One thousand five hundred and thirty cases out of 3670 (44.1%, 95% CI 39.5% to 48.9%) cases had radiological abnormalities; ground glass opacities (27.4%) were the most commonly reported abnormality.

Sixty-six studies provided details on laboratory-markers (table 2). Pooled analysis revealed increased C-Reactive Protein (CRP) (54.2%, 95% CI 41.5% to 66.3%), serum-ferritin (46.7%, 95% CI 32.3% to 61.7%), lactate dehydrogenase (LDH) (36.5%, 95% CI 26.5% to 47.8%) and d-dimers (35.2%, 95% CI 22.1% to 51.0%) as the most common abnormalities. Other reported abnormalities included elevated erythrocyte sedimentation rate (ESR), lymphopaenia, procalcitonin and biomarkers for organ

**Table 1** Clinical symptoms among reported paediatric COVID-19 cases

Characteristics	Events/total patients	Mean proportion % (95% CI)	Heterogeneity I <sup>2</sup> (%)
Comorbidity	1590/6086	27.1 (23.1 to 31.5)	37.6
Fever	3576/6296	63.3 (58.6 to 68.4)	34.9
Cough	1807/5261	33.7 (29.6 to 38.1)	34.4
Nausea/vomiting	880/4243	20.0 (16.5 to 24.0)	25.7
Diarrhoea	796/4884	19.6 (16.1 to 23.7)	13.4
Dyspnoea	879/5332	17.5 (14.4 to 21.1)	23.7
Nasal symptoms	1080/5406	16.6 (13.9 to 19.7)	10.6
Rashes	744/4387	15.5 (11.9 to 19.9)	25.9
Fatigue	709/4474	15.5 (12.6 to 19.3)	26.3
Abdominal pain	626/4135	15.3 (11.9 to 19.4)	26.5
Kawasaki shock/sign	821/4365	13.3 (9.8 to 17.9)	30.6
Asymptomatic	1114/6084	13.1 (10.4 to 16.3)	15.4
Neurological symptoms	693/5475	12.1 (10.1 to 14.6)	17.6
Conjunctivitis	529/4998	10.5 (7.8 to 14.0)	21.0
Pharyngeal erythema	428/3638	9.0 (6.7 to 12.0)	0.0

injury including elevated levels of pro B-type natriuretic peptide, troponin and creatine kinase-MB as shown in [table 2](#).

### MANAGEMENT

Details of clinical management are as shown in [table 3](#). Commonly used therapies among hospitalised children were antimicrobials (32.2%, 95% CI 25.2% to 40.1%), intravenous immunoglobulin (IVIG) (19.5%, 95% CI 13.5% to 27.2%) and systemic-steroids (19.3, 95% CI 14.9% to 24.9%). Other treatment regimens included aspirin, inotropic drugs, inhaled interferon- $\alpha$  (IFN- $\alpha$ ), antimalarials and antivirals (ribavirin, oseltamivir, lopinavir,

ritonavir and litonavir). Mechanical ventilation was provided to 490 patients (12.2%, 95% CI 9.7% to 15.3%).

### PROGNOSIS AND SEVERE CASES

One thousand three hundred and fifty-nine patients (22.9%, 95% CI 17.6% to 29.2%) were admitted to ICUs ([table 3](#)). Thirty-eight studies provided disaggregated data for severe cases ([table 4](#)). A higher proportion of children with severe disease had symptoms consistent with MIS-C and received antimicrobials, inotropes and anti-inflammatory agents compared with those with non-severe disease. There were no deaths among children

**Table 2** Laboratory and radiological features among reported paediatric COVID-19 cases

Characteristics	Events/total patients	Mean proportion % (95% CI)	Heterogeneity I <sup>2</sup> (%)
<b>Inflammatory marker</b>			
CRP $\uparrow$	556/1165	54.2 (41.5 to 66.3)	21.4
Ferritin $\uparrow$	247/525	46.7 (32.3 to 61.7)	46.5
LDH $\uparrow$	356/922	36.5 (26.5 to 47.8)	35.6
Procalcitonin $\uparrow$	137/879	21.3 (12.2 to 34.5)	24.9
Leukocytes $\uparrow$	138/953	19.9 (13.3 to 28.8)	21.4
Lymphocytes $\downarrow$	359/1347	19.0 (12.8 to 27.1)	0.0
ESR $\uparrow$	248/838	18.9 (11.8 to 28.9)	0.0
IL-6 $\uparrow$	41/341	13.1 (5.5 to 28.2)	7.1
Leucopaenia (+)	77/1037	10.7 (7.7 to 14.6)	0.0
Lymphocytes $\uparrow$	66/1264	8.2 (4.9 to 13.5)	0.0
Neutrophils $\uparrow$	22/574	7.8 (4.8 to 12.4)	0.0
<b>Biomarkers for organ injury</b>			
proBNP $\uparrow$	211/441	45.5 (28.5 to 63.5)	49.5
Troponin $\uparrow$	239/703	39.7 (24.7 to 57.0)	30.5
LFTs $\uparrow$	287/816	29.8 (20.3 to 41.6)	10.8
CKMB $\uparrow$	82/293	25.5 (13.4 to 43.0)	31.1
RFTs $\uparrow$	86/344	17.6 (7.6 to 35.6)	23.6
<b>Coagulopathy markers</b>			
D-dimers $\uparrow$	272/711	35.2 (22.1 to 51.0)	19.1
Fibrinogen $\uparrow$	168/438	17.5 (7.6 to 35.4)	0.0
<b>Radiological test</b>			
Abnormal CXR/CT	1530/3670	44.1 (39.5 to 48.9)	35.0

**Table 3** Clinical management and outcomes among reported paediatric COVID-19 cases

Characteristics	Events/total patients	Mean proportion % (95% CI)	Heterogeneity I <sup>2</sup> (%)
<b>Clinical management</b>			
Antibiotics	1345/3610	32.2 (25.2 to 40.1)	41.9
IVIg	698/3522	19.5 (13.5 to 27.2)	18.4
Systemic steroids	801/4229	19.3 (14.9 to 24.9)	23.7
Antiviral	527/4019	15.3 (11.1 to 20.7)	4.5
Mechanical ventilation	490/5406	12.2 (9.7 to 15.3)	15.5
Inotropes	354/3856	11.8 (8.3 to 16.4)	11.5
Antimalarial	336/3299	9.9 (6.9 to 14.0)	0.0
Aspirin	238/2588	9.0 (5.9 to 13.6)	78.1
Interferon	138/2598	7.7 (4.9 to 11.8)	0.0
Traditional medicine	22/4229	4.0 (2.8 to 5.6)	38.7
<b>Clinical outcomes</b>			
Recovered	8704/9335	88.9 (86.0 to 91.2)	36.3
ICU admission	1359/9335	22.9 (17.6 to 29.2)	37.2
Deaths	96/6902	3.6 (2.8 to 4.5)	24.3

ICU, intensive care unit; IVIG, intravenous immunoglobulin.

**Table 4** Comparison of clinical symptoms, management and outcomes among reported paediatric COVID-19 non-severe (n=2402 cases, 64 studies) and severe (n=796 cases, 38 studies) cases

Characteristics	Non-severe cases		Severe cases		RR severe vs non-severe (95% CI)
	Events/total patients	Mean proportion % (95% CI)	Events/total patients	Mean proportion % (95% CI)	
<b>Clinical symptoms</b>					
Fever	1394/2404	51.4 (45.7 to 57.0)	608/756	80.2 (73.6 to 85.5)	1.39 (1.32 to 1.46)
Pharyngeal erythema	541/1149	8.6 (5.1 to 14.0)	41/585	8.8 (5.1 to 14.8)	0.15 (0.11 to 0.20)
Cough	587/1521	35.1 (29.2 to 41.5)	225/618	34.0 (24.6 to 44.9)	0.94 (0.84 to 1.07)
Comorbidity	541/2283	19.8 (14.5 to 26.4)	351/764	44.1 (34.9 to 53.8)	1.94 (1.74 to 2.16)
Nausea/vomiting	206/1291	12.1 (8.7 to 16.6)	224/632	41.0 (36.7 to 45.5)	2.27 (1.93 to 2.67)
Dyspnoea	260/1646	12.7 (9.5 to 16.8)	237/701	36.4 (26.5 to 47.5)	2.14 (1.84 to 2.49)
Nasal symptoms	402/1659	14.1 (9.9 to 19.7)	91/652	15.8 (10.6 to 23.0)	0.58 (0.47 to 0.71)
Fatigue	192/1319	13.8 (10.4 to 18.0)	151/505	20.3 (15.3 to 33.4)	2.05 (1.70 to 2.48)
Kawasaki shock/sign	135/1243	8.5 (5.6 to 12.6)	242/695	30.7 (19.3 to 45.0)	3.21 (2.65 to 3.87)
Rashes	168/1587	10.3 (7.6 to 13.7)	180/660	32.3 (22.3 to 44.2)	2.58 (2.13 to 3.11)
Abdominal pain	95/1193	8.1 (5.8 to 11.3)	184/621	28.4 (18.8 to 40.4)	3.72 (2.96 to 4.67)
Diarrhoea	144/1326	13.5 (10.6 to 17.1)	217/632	35.3 (26.3 to 45.4)	3.16 (2.62 to 3.82)
Conjunctivitis	111/1621	7.5 (5.1 to 10.8)	116/657	22.6 (15.1 to 32.4)	2.58 (2.02 to 3.29)
Neurological symptoms	200/2230	11.0 (9.0 to 13.4)	118/703	17.4 (11.9 to 24.6)	1.87 (1.52 to 2.31)
<b>Clinical management</b>					
Mechanical ventilation	–	–	322/735	43.8 (33.8 to 54.3)	–
Antiviral	217/715	26.5 (17.5 to 38.1)	136/567	24.1 (16.2 to 34.3)	0.79 (0.66 to 0.95)
Interferon	127/685	20.2 (11.6 to 32.3)	4/445	6.8 (3.6 to 12.3)	0.05 (0.02 to 0.13)
Antibiotics	180/363	21.6 (14.2 to 31.3)	365/566	59.6 (44.3 to 73.3)	1.30 (1.15 to 1.47)
Antimalarial	73/717	10.1 (6.4 to 16.4)	123/537	22.9 (14.3 to 34.6)	2.25 (1.72 to 2.94)
IVIg	54/721	11.8 (7.2 to 18.8)	202/498	41.1 (27.0 to 56.8)	5.42 (4.10 to 7.15)
Systemic steroids	46/721	8.9 (5.4 to 14.4)	265/575	46.8 (35.7 to 58.2)	7.22 (5.39 to 9.69)
Inotropes	24/718	6.8 (4.1 to 11.7)	171/498	33.6 (21.2 to 48.9)	10.27 (6.81 to 15.50)
Traditional medicine	18/723	7.4 (4.8 to 11.3)	4/575	6.3 (3.8 to 10.3)	0.28 (0.10 to 0.82)
Aspirin	11/683	6.7 (4.3 to 10.4)	83/445	14.9 (7.9 to 26.4)	11.58 (6.25 to 21.47)
<b>Clinical outcomes</b>					
ICU	–	–	793/796	95.0 (92.1 to 96.8)	–
Recovered	1700/1925	85.4 (76.5 to 91.2)	532/796	77.6 (67.5 to 85.3)	0.76 (0.72 to 0.80)
Deaths	0/1925	–	44/796	8.0 (5.2 to 12.1)	–

ICU, intensive care unit; IVIG, intravenous immunoglobulin.

**Table 5** Comparison of clinical symptoms, management and outcomes among reported paediatric COVID-19 cases in HICs (n=5641 cases, 60 studies) and LMICs (n=3694, 69 studies)

Characteristics	HICs		LMICs		RR LMICs vs HICs, (95% CI)
	Events/total patients	Mean proportion % (95% CI)	Events/total patients	Mean proportion % (95% CI)	
<b>Clinical symptoms</b>					
Fever	2276/3332	72.0 (66.3 to 77.0)	1300/2964	50.0 (47.4 to 52.6)	0.64 (0.61 to 0.67)
Cough	995/2730	33.2 (27.5 to 39.5)	812/2531	39.2 (36.2 to 42.3)	0.88 (0.82 to 0.95)
Comorbidity	1069/3357	33.7 (27.4 to 38.5)	521/2729	20.8 (18.3 to 23.4)	0.60 (0.55 to 0.66)
Rashes	491/2109	24.9 (17.9 to 33.5)	253/2278	20.2 (17.5 to 23.2)	0.48 (0.41 to 0.55)
Nausea/vomiting	668/2374	30.3 (24.3 to 37.0)	212/1869	15.3 (12.6 to 18.3)	0.40 (0.35 to 0.46)
Conjunctivitis	309/2732	13.1 (8.4 to 20.1)	220/2266	19.5 (16.9 to 22.5)	0.86 (0.73 to 1.01)
Dyspnoea	543/2454	23.6 (18.5 to 29.6)	336/2878	20.6 (18.1 to 23.4)	0.53 (0.47 to 0.60)
Kawasaki shock/sign	583/2087	21.9 (14.1 to 32.5)	238/2278	21.4 (18.5 to 24.6)	0.37 (0.33 to 0.43)
Fatigue	394/1943	16.8 (12.2 to 22.6)	315/2531	15.6 (13.3 to 18.3)	0.61 (0.54 to 0.70)
Abdominal pain	457/2266	22.7 (16.4 to 30.5)	169/1869	16.6 (12.9 to 21.1)	0.45 (0.38 to 0.53)
Nasal symptoms	425/2549	17.8 (14.4 to 21.8)	269/2519	15.4 (13.2 to 17.8)	0.64 (0.56 to 0.74)
Diarrhoea	527/2365	27.5 (21.5 to 34.6)	125/1105	14.7 (12.4 to 17.3)	0.51 (0.42 to 0.61)
Neurological symptoms	493/3197	15.0 (11.6 to 19.0)	200/2278	10.4 (8.8 to 12.3)	0.57 (0.49 to 0.67)
Asymptomatic	263/2428	6.4 (4.2 to 9.7)	851/3656	20.2 (18.4 to 22.1)	2.15 (1.89 to 2.44)
Pharyngeal erythema	73/1494	6.7 (4.3 to 10.1)	519/2531	40.7 (37.4 to 44.0)	4.20 (3.31 to 5.32)
<b>Clinical management</b>					
Antibiotics	908/1875	36.4 (25.6 to 48.7)	437/1735	27.0 (22.9 to 31.5)	0.52 (0.47 to 0.57)
IVIG	504/1867	31.6 (20.3 to 45.5)	194/1655	14.7 (12.1 to 17.7)	0.43 (0.37 to 0.51)
Aspirin	187/985	16.0 (9.1 to 26.8)	51/1603	10.4 (7.2 to 14.7)	0.17 (0.12 to 0.23)
Systemic steroids	566/2523	27.2 (19.0 to 37.3)	235/1706	18.0 (15.2 to 21.2)	0.61 (0.53 to 0.71)
Inotropes	309/2309	19.1 (12.3 to 28.5)	45/1547	11.9 (8.5 to 16.5)	0.22 (0.16 to 0.30)
Antimalarial	241/1696	13.6 (8.7 to 20.8)	95/1603	13.5 (10.5 to 17.3)	0.42 (0.33 to 0.52)
Mechanical ventilation	387/2930	17.2 (13.1 to 22.3)	103/2476	10.8 (8.6 to 13.4)	0.32 (0.26 to 0.39)
Antiviral	230/2372	11.4 (7.7 to 16.6)	297/1647	25.2 (21.1 to 29.9)	1.86 (1.58 to 2.18)
Interferon	0/995	–	138/1603	30.5 (24.1 to 37.7)	–
Traditional medicine	0/2523	–	22/1706	11.3 (8.0 to 15.7)	–
<b>Clinical outcomes</b>					
Recovered	5269/5641	91.0 (87.7 to 93.4)	3435/3694	83.9 (81.2 to 86.2)	0.99 (0.98 to 1.01)
ICU admission	993/5641	26.0 (24.0 to 28.0)	366/3694	9.9 (8.5 to 11.6)	0.56 (0.50 to 0.63)
Deaths	40/4710	2.9 (2.1 to 4.1)	56/2192	5.2 (4.1 to 6.7)	2.14 (1.43 to 3.20)

HICs, high-income countries; ICU, intensive care unit; IVIG, intravenous immunoglobulin; LMICs, low-income and middle-income countries; RR, relative risk.

categorised as non-severe and 44 deaths among severe cases, where disaggregated data were provided. Hospital outcomes were reported for 9335 children; 8704 cases (88.9%, 95% CI 86.0% to 91.2%) were definitively discharged, 96 died and remaining children either remained hospitalised at the time of reporting or were readmitted.

### COMPARISON OF OUTCOMES ACCORDING TO WORLD BANK COUNTRY CLASSIFICATION

Of the included studies that reported clinical characteristics and outcomes, 60 studies were from HICs (n=6528) and 69 studies from LMICs (n=3723) as show in table 5. Studies in LMICs included a higher proportion of hospitalised children (1981/3723, 53.2%) compared with HIC studies (2897/6528, 44.4%). Abdominal symptoms and symptoms consistent with MIS-C were more frequently reported in HICs. A lower proportion of children in LMICs were admitted to the ICU (RR 0.56, 95% CI 0.50 to 0.63,  $p<0.05$ ), mechanically ventilated (RR 0.32, 95% CI 0.26 to 0.39,  $p<0.05$ ) and treated with different

therapies; inotropes, antimicrobials, steroids, aspirin and IVIG. Only children in LMICs received inhaled IFN- $\alpha$ . Among the hospitalised cases, 40 deaths were reported in HICs compared with 56 in LMICs (pooled proportion 2.9% vs 5.2%). Risk-adjusted mortality according to severity of illness could not be calculated due to lack of data (table 5).

### SUBGROUP ANALYSES OF CHILDREN PRESENTING WITH MIS-C, AND COVID-19 IN NEONATES

Thirty-one studies (n=1208) with 22 from HIC (n=602), reported series of children presenting with MIS-C. Fever, abdominal pain and diarrhoea were the most common symptoms. Nearly half of children (638/1208) who met criteria for MIS-C were admitted to ICU (449/638, 70.3% of which were from HIC) compared with 22.9% in the overall analysis (online supplemental tables 6 and 7).

Disaggregated data were available on 184 neonates with fever; inability to feed/lethargy and dyspnoea were the most commonly

reported symptoms. Twenty-one neonates (16.6%, 95% CI 11.2% to 23.9%) were asymptomatic at the time of diagnosis.

### QUALITY ASSESSMENT OF INCLUDED STUDIES

One hundred and twenty-one studies were determined to be of good quality while eight were of fair quality (online supplemental table 8). Studies were primarily downgraded for incomplete case definition,<sup>29 31 44 46 48–50 130 135 138</sup> incomplete case follow-up,<sup>10 23 24 26 29 32 35 44 51 53 77 82 85 90 94 99 106 112–114 120 130 135–137</sup> missing data<sup>2 3 19–21 29 35 42–44 52 57 74 89 92–94 117 128</sup> and non-consecutive patient enrolment,<sup>9 11 12 16 18–26 28 30–35 37–39 50 57 61–73 80 82 87 89–91 93 95 102 109 110 112 114 123 125 126 129 134–136</sup> which raises concern that the included sample could be biased towards more severe presentations.

### DISCUSSION

Global knowledge of COVID-19 epidemiology, clinical characteristics and management has continued to evolve since the onset of the pandemic. Children have been noted to have relatively lower rates of severe illness and low mortality; however, they have been impacted by MIS-C.<sup>4 139</sup>

The findings of our review, the largest in terms of published systematic reviews on paediatric COVID-19, are consistent with previous reviews that identified predominance of infection in school-age children, slight male predisposition, prevalence of comorbidities among children with COVID-19 and low hospitalisation and mortality rates.<sup>2 4 140</sup> The clinical presentation in children is heterogeneous, including a wide spectrum of clinical features. Fever and cough were the most commonly reported presenting symptoms, in line with the previously published systematic reviews.<sup>4 141</sup> A U-shaped curve of severity has been demonstrated in children diagnosed with COVID-19 with infants under 1 year of age and adolescents 10–14 years of age at higher risk of developing severe COVID-19.<sup>3 47 75</sup> Due to lack of age-disaggregated data, we could not reliably compare the frequency of severe cases by age group in this review. Reported risk factors for severe disease among children include age, viral load<sup>142</sup> and presence of comorbidities.<sup>3</sup> There is a possibility that children with comorbidities may have been hospitalised related to their underlying chronic condition and incidentally determined to have COVID-19 infection or investigated more extensively. Some of the common comorbidities reported in children with COVID-19 infection include asthma, immunosuppression, congenital heart disease, kidney disease and obesity.<sup>3 4 47</sup>

Regional differences were identified in the comparison of clinical features, treatment and outcomes between HICs and LMICs. Pooled estimates of hospital mortality were higher in LMICs compared with HICs. Given that it was not possible to calculate risk-adjusted mortality rates for COVID-19, it is unclear whether observed differences in mortality are related to selection bias (eg, differences in severity of illness of included patients or differences in case definitions and inclusion criteria) or differences in available hospital resources. Nevertheless, there is ongoing concern that, in LMICs with high burden of illness and health system limitations, children with severe disease and MIS-C may be at greater risk for adverse outcomes and death than perceived to date. The differences in frequency of observed clinical features may be related to increasing recognition of MIS-C over the course of the pandemic and their inclusion in more recent COVID-19 case series, but is likely similar between HICs and LMICs.

Comparisons of clinical features and outcomes according to severity of illness were limited by heterogeneous reporting

across the included case series. A higher proportion of children with severe disease demonstrated symptoms consistent with MIS-C (fever, abdominal symptoms, rash, neurological symptoms, conjunctivitis) and received IVIG, steroids and inotropes.

Compared with previous reviews, several at an earlier stage of the pandemic,<sup>4 7 8 140 143</sup> this review has several strengths. Using a broad search strategy implemented in English, Chinese and Spanish databases, we summarise evidence from 129 studies from 31 different countries, the largest sample to-date. We excluded case reports to minimise selective reporting of extreme and atypical cases. We also attempted to reduce possible overlap in cases to prevent duplications. We identified differences in features from studies in HICs compared with LMICs, and between severe and non-severe cases, although with limited available data. Finally, we report subgroup analyses for neonates, and children presenting with MIS-C.

The review is limited primarily by the small sample sizes of individual studies, limitations in study reporting, and study quality limitations due to non-consecutive patient enrolment, unclear case definition and incomplete follow-up to hospital discharge. Our approach of pooling proportions is subject to bias and wide confidence-intervals due to small study sample size. We could not undertake multivariate analysis to identify risk factors for severe infection or adverse outcome in children due to lack of individual-patient-data. The inclusion of asymptomatic cases could have contributed to underestimation of the prevalence of clinical characteristics and optimism in the reporting of outcomes. Finally, it should be noted that a large number (36/129, 28.0%) of the included studies were from China. While the Chinese healthcare system is well-resourced in certain regions, many of the Chinese studies included were conducted in the city of Wuhan or in Hubei Province (n=9, 32.1%), where the gross domestic product per capita is less than half of that of Beijing and Shanghai.<sup>144</sup> Therefore, the findings of studies from China may be generalisable to the socioeconomic and health development status of other middle-income countries.

This review contributes to the global understanding of paediatric COVID-19 disease and supports priority setting in research for current pandemic and future outbreaks. This body of literature would be improved by complete reporting of larger series with consecutive recruitment of patients, specific case definitions and complete long-term follow-up to determine global epidemiological trends, age-specific burden of disease and illness trajectory following COVID-19 infection. Improved characterisation of disease severity and increased reports from low-income countries are needed to better understand differences in clinical manifestations, resource utilisation and outcome by region, which can be integrated in future updated analyses. The concern for selection bias remains as it is possible that in LMICs, the population of hospitalised children was sicker and at higher baseline risk of death, independent of resources. Individual-patient-data meta-analysis would be of benefit to characterise risk factors for severe disease, clinical features in different age groups and account for observed differences in outcome. With respect to clinical management, none of the therapies instituted in the treatment of children with severe COVID-19 disease have been demonstrated to improve outcome in randomised trials; therefore, a recommendation regarding their use is challenging. Given that children appear less likely to develop severe respiratory disease, but are at risk of multiorgan dysfunction due to MIS-C, further studies are needed to characterise the clinical trajectory of this novel syndrome and determine the optimal treatment for it. Finally, there remains paucity of studies reporting long-term prognosis of COVID-19 in children.<sup>145</sup>

## CONCLUSION

Our review suggests that children predominantly contracted mild form of infection but could be at risk of more severe outcomes. It is crucial to take into consideration risk factors including contact-exposure, underlying comorbidities, young age and male sex which may increase the risk of severe disease. While we have identified several elements that highlight the disease spectrum and higher risk of adverse outcomes in certain settings, such as LMICs, there is the need for much closer scrutiny of this illness globally with individual patient data analysis.

**Contributors** ZB conceptualised the study and secured funding. ZSL and OI drafted the study protocol, conducted the literature search, study screening, selection and data extraction and drafted the manuscript. LJ and KT designed the data collection instruments, collected data, carried out data analyses and reviewed and revised the manuscript. FM drafted the initial manuscript and reviewed and revised the manuscript. All authors critically reviewed the manuscript for important intellectual content and approved the final manuscript as submitted. ZB is the guarantor.

**Funding** This review was funded in part by a grant from UNICEF (Headquarters) in partnership with the International Pediatric Association and with core support from the Centre for Global Child Health (Toronto) and the Center of Excellence in Women & Child Health, The Aga Khan University, Karachi, Pakistan

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; internally peer reviewed.

**Data availability statement** Data are available upon request

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iD

Zulfiqar Bhutta <http://orcid.org/0000-0003-0637-599X>

## REFERENCES

- WHO. Who coronavirus disease (COVID-19) Dashboard data last updated: 2021/1/21. Available: <https://covid19.who.int/>
- Dong Y, Mo X, Hu Y. Epidemiological characteristics of 2143 pediatric patients with 2019 coronavirus disease in China. *J Emerg Med* 2020;58:712–3.
- Covid CD CC, COVID C, Bialek S, et al. Coronavirus Disease 2019 in Children - United States, February 12–April 2, 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:422.
- Hoang A, Chorath K, Moreira A, et al. COVID-19 in 7780 pediatric patients: a systematic review. *EclinicalMedicine* 2020;24:100433.
- AAP. *Children and COVID19: state data report 28/8/2020*, 2020.
- Wang E, Brar K. COVID-19 in children: an epidemiology study from China. *J Allergy Clin Immunol* 2020;8:2118–20.
- de Souza TH, Nadal JA, Nogueira RJN, et al. Clinical manifestations of children with COVID-19: a systematic review. *Pediatr Pulmonol* 2020;55:1892–9.
- Castagnoli R, Votto M, Licari A, et al. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection in children and adolescents. *JAMA Pediatr* 2020;174:882.
- Cheung EW, Zachariah P, Gorelik M, et al. Multisystem inflammatory syndrome related to COVID-19 in previously healthy children and adolescents in New York City. *JAMA* 2020;324:294.
- Feldstein LR, Rose EB, Horwitz SM, et al. Multisystem inflammatory syndrome in U.S. children and adolescents. *N Engl J Med Overseas Ed* 2020;383:334–46.
- Blondiaux E, Parisot P, Redheuil A, et al. Cardiac MRI in children with multisystem inflammatory syndrome associated with COVID-19. *Radiology* 2020;297:E283–8.
- Pouletty M, Borocco C, Ouldali N, et al. Paediatric multisystem inflammatory syndrome temporally associated with SARS-CoV-2 mimicking Kawasaki disease (Kawa-COVID-19): a multicentre cohort. *Ann Rheum Dis* 2020;79:999–1006.
- Global Financing Facility for Women, Children and Adolescents (GFF)- September 18, 2020. Available: <https://www.globalfinancingfacility.org/new-findings-confirm-global-disruptions-essential-health-services-women-and-children-covid-19>
- National Heart L, and Blood Institute. *Quality assessment tool for observational cohort and cross-sectional studies*. Bethesda: National Institutes of health, department of health and human services, 2014.
- Group WB. *World bank country and lending groups (country classification)*, 2020.
- Qiu H, Wu J, Hong L, et al. Clinical and epidemiological features of 36 children with coronavirus disease 2019 (COVID-19) in Zhejiang, China: an observational cohort study. *Lancet Infect Dis* 2020;20:689–96.
- Wang D, Ju XL, Xie F, et al. [Clinical analysis of 31 cases of 2019 novel coronavirus infection in children from six provinces (autonomous region) of northern China]. *Zhonghua Er Ke Za Zhi* 2020;58:E011–E.
- Zheng F, Liao C, Fan Q-H, et al. Clinical characteristics of children with coronavirus disease 2019 in Hubei, China. *Curr Med Sci* 2020;40:275–80.
- Xia W, Shao J, Guo Y, et al. Clinical and CT features in pediatric patients with COVID-19 infection: different points from adults. *Pediatr Pulmonol* 2020;55:1169–74.
- Feng K, Yun YX, Wang XF, et al. [Analysis of CT features of 15 Children with 2019 novel coronavirus infection]. *Zhonghua Er Ke Za Zhi* 2020;58:E007.
- Jiehao C, Jin X, Daojiong L, et al. A case series of children with 2019 novel coronavirus infection: clinical and epidemiological features. *Clin Infect Dis* 2020;71:1547–51.
- Wei M, Yuan J, Liu Y, et al. Novel coronavirus infection in hospitalized infants under 1 year of age in China. *JAMA* 2020;323:1313–4.
- Su L, Ma X, Yu H, et al. The different clinical characteristics of corona virus disease cases between children and their families in China - the character of children with COVID-19. *Emerg Microbes Infect* 2020;9:707–13.
- Zhou Y, Yang G-D, Feng K, et al. [Clinical features and chest CT findings of coronavirus disease 2019 in infants and young children]. *Zhongguo Dang Dai Er Ke Za Zhi* 2020;22:215–20.
- Sun D, Li H, Lu X-X, et al. Clinical features of severe pediatric patients with coronavirus disease 2019 in Wuhan: a single center's observational study. *World J Pediatr* 2020;16:251–9.
- Liu W, Zhang Q, Chen J, et al. Detection of Covid-19 in children in early January 2020 in Wuhan, China. *N Engl J Med* 2020;382:1370–1.
- Hu Z, Song C, Xu C, et al. Clinical characteristics of 24 asymptomatic infections with COVID-19 screened among close contacts in Nanjing, China. *Sci China Life Sci* 2020;63:706–11.
- Liu M, Song Z, Xiao K. High-Resolution computed tomography manifestations of 5 pediatric patients with 2019 novel coronavirus. *J Comput Assist Tomogr* 2020;44:311–3.
- Liu H, Liu F, Li J, et al. Clinical and CT imaging features of the COVID-19 pneumonia: focus on pregnant women and children. *J Infect* 2020;80:e7–e13.
- Lou XX, Shi CX, Zhou CC, et al. Three children who recovered from novel coronavirus 2019 pneumonia. *J Paediatr Child Health* 2020;56:650–651.
- Zhang T, Cui X, Zhao X, et al. Detectable SARS-CoV-2 viral RNA in feces of three children during recovery period of COVID-19 pneumonia. *J Med Virol* 2020;92:909–14.
- Li Y, Guo F, Cao Y, et al. Insight into COVID-2019 for pediatricians. *Pediatr Pulmonol* 2020;55:E1–4.
- Ji L-N, Chao S, Wang Y-J, et al. Clinical features of pediatric patients with COVID-19: a report of two family cluster cases. *World J Pediatr* 2020;16:267–70.
- Tagarro A, Epalza C, Santos M, et al. Screening and severity of coronavirus disease 2019 (COVID-19) in children in Madrid, Spain. *JAMA Pediatr* 2020. doi:10.1001/jamapediatrics.2020.1346. [Epub ahead of print: 08 Apr 2020].
- Chen A, Huang J-xiang, Liao Y, et al. Differences in clinical and imaging presentation of pediatric patients with COVID-19 in comparison with adults. *Radiology* 2020;2:e200117.
- See KC, Liew SM, Ng DCE, et al. COVID-19: four paediatric cases in Malaysia. *Int J Infect Dis* 2020;94:125–7.
- Zhang G-X, Zhang A-M, Huang L, et al. [Twin girls infected with SARS-CoV-2]. *Zhongguo Dang Dai Er Ke Za Zhi* 2020;22:221–5.
- Cheng. *A case report of two children with corona virus disease 2019 complicated with diffuse intravascular coagulation*, 2020.
- Tang A, Xu W, Chen P. A retrospective study of the clinical characteristics of COVID-19 infection in 26 children. *medRxiv* 2020.
- Zhu L, Wang J, Huang R, et al. Clinical characteristics of a case series of children with coronavirus disease 2019. *Pediatr Pulmonol* 2020;55:1430–2.
- Tan X, Huang J, Zhao F, et al. [Clinical features of children with SARS-CoV-2 infection: an analysis of 13 cases from Changsha, China]. *Zhongguo Dang Dai Er Ke Za Zhi* 2020;22:294.
- Ma H, Shao J, Wang Y. High resolution CT features of COVID-19 in children. *Chinese Journal of Radiology* 2020;54.
- Shen Q, Guo W, Guo T, et al. Novel coronavirus infection in children outside of Wuhan, China. *Pediatr Pulmonol* 2020;55:1424–9.

- 44 Schwierzeck V, König JC, Kühn J, *et al.* First reported nosocomial outbreak of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in a pediatric dialysis unit. *Clin Infect Dis* 2020. doi:10.1093/cid/ciaa491. [Epub ahead of print: 27 Apr 2020].
- 45 Jiang J, Duan L, Xiong D. Epidemiological and clinical characteristics of novel coronavirus infection in children: thoughts on the diagnostic criteria of suspected cases outside Hubei Province. *Chinese Pediatric Emergency Medicine* 2020.
- 46 Riphagen S, Gomez X, Gonzalez-Martinez C, *et al.* Hyperinflammatory shock in children during COVID-19 pandemic. *The Lancet* 2020;395:1607–8.
- 47 Shekerdemian LS, Mahmood NR, Wolfe KK, *et al.* Characteristics and outcomes of children with coronavirus disease 2019 (COVID-19) infection admitted to US and Canadian pediatric intensive care units. *JAMA Pediatr* 2020;174:868.
- 48 Verdoni L, Mazza A, Gervasoni A, *et al.* An outbreak of severe Kawasaki-like disease at the Italian epicentre of the SARS-CoV-2 epidemic: an observational cohort study. *The Lancet* 2020;395:1771–8.
- 49 Rahimzadeh G, Ekrami Noghahi M, Kadkhodaei Elyaderani F, *et al.* COVID-19 infection in Iranian children: a case series of 9 patients. *J Pediatr Rev* 2020;8:139–44.
- 50 Tullie L, Ford K, Bisharat M, *et al.* Gastrointestinal features in children with COVID-19: an observation of varied presentation in eight children. *Lancet Child Adolesc Health* 2020;4:e19–20.
- 51 Sajid MI, Altaf S, Mushtaq N. A wrinkle in time: how has management of pediatric cancers changed during COVID-19 in Pakistan? 2020.
- 52 Zhang Z-J, Yu X-J, Fu T, *et al.* Novel coronavirus infection in newborn babies aged <28 days in China. *Eur Respir J* 2020;55:2000697.
- 53 Zeng L, Xia S, Yuan W, *et al.* Neonatal early-onset infection with SARS-CoV-2 in 33 neonates born to mothers with COVID-19 in Wuhan, China. *JAMA Pediatr* 2020;174:722.
- 54 Salvatori G, De Rose DU, Concato C, *et al.* Managing COVID-19-Positive maternal-infant dyads: an Italian experience. *Breastfeed Med* 2020;15:347–8.
- 55 Zhang XMY, Xiao J, Zhang Z. Clinical characteristics of novel coronavirus pneumonia in children in Jinan.
- 56 Wu H-P, Li B-F, Chen X, *et al.* [Clinical features of coronavirus disease 2019 in children aged <18 years in Jiangxi, China: an analysis of 23 cases]. *Zhongguo Dang Dai Er Ke Za Zhi* 2020;22:419–24.
- 57 Zhang H-Y, Zhang Y. [Coronavirus disease 2019 and hypertension in 2 children]. *Zhongguo Dang Dai Er Ke Za Zhi* 2020;22:425–8.
- 58 Chao JY, Derespina KR, Herold BC, *et al.* Clinical characteristics and outcomes of hospitalized and critically ill children and adolescents with coronavirus disease 2019 at a tertiary care medical center in New York City. *J Pediatr* 2020;223:14–19.
- 59 Eghbali ASS, Sadat N, *et al.* COVID-19 in pediatric patients: a case series. *J Cell Mol Med* 2020;5:3–5.
- 60 Chiotos K, Bassiri H, Behrens EM, *et al.* Multisystem inflammatory syndrome in children during the coronavirus 2019 pandemic: a case series. *J Pediatric Infect Dis Soc* 2020;9:393–8.
- 61 Flores V, Miranda R, Merino L, *et al.* SARS-CoV-2 infection in children with febrile neutropenia. *Ann Hematol* 2020;99:1–2.
- 62 García-Salido A, Leoz-Gordillo I, Martínez de Azagra-Garde A, *et al.* Children in critical care due to severe acute respiratory syndrome coronavirus 2 infection: experience in a Spanish hospital. *Pediatr Crit Care Med* 2020;21:e576–e580.
- 63 Cabrero-Hernández M, García-Salido A, Leoz-Gordillo I, *et al.* Severe SARS-CoV-2 infection in children with suspected acute abdomen: a case series from a tertiary hospital in Spain. *Pediatr Infect Dis J* 2020;39:e195–8.
- 64 Waltuch T, Gill P, Zinns LE, *et al.* Features of COVID-19 post-infectious cytokine release syndrome in children presenting to the emergency department. *Am J Emerg Med* 2020;38:2246.e3–2246.e6.
- 65 Kakuya F, Okubo H, Fujiyasu H, *et al.* The first pediatric patients with coronavirus disease 2019 (COVID-19) in Japan: risk of co-infection with other respiratory viruses. *Jpn J Infect Dis* 2020;73:377–80.
- 66 Soltani J, Sedighi I, Shalchi Z, *et al.* Pediatric coronavirus disease 2019 (COVID-19): an insight from West of Iran. *North Clin Istanb* 2020;7:284.
- 67 Toubiana J, Poirault C, Corsia A, *et al.* Kawasaki-like multisystem inflammatory syndrome in children during the covid-19 pandemic in Paris, France: prospective observational study. *BMJ* 2020;369:m2094.
- 68 Foster CE, Moulton EA, Munoz FM, *et al.* Coronavirus disease 2019 in children Cared for at Texas children's Hospital: initial clinical characteristics and outcomes. *J Pediatric Infect Dis Soc* 2020;9:373–7.
- 69 White A, Mukherjee P, Stremming J, *et al.* Neonates hospitalized with community-acquired SARS-CoV-2 in a Colorado neonatal intensive care unit. *Neonatology* 2021;117:641–5.
- 70 Parri N, Lenge M, Cantoni B, *et al.* COVID-19 in 17 Italian pediatric emergency departments. *Pediatrics* 2020;146. doi:10.1542/peds.2020-1235. [Epub ahead of print: 23 Sep 2020].
- 71 Gao Y, Zhang D, Sui S, *et al.* Clinical features and treatment protocol in eleven Chinese children with mild COVID-19. *Indian J Pediatr* 2020;87:748.
- 72 Harman K, Verma A, Cook J, *et al.* Ethnicity and COVID-19 in children with comorbidities. *Lancet Child Adolesc Health* 2020;4:e24–5.
- 73 Whittaker E, Bamford A, Kenny J, *et al.* Clinical characteristics of 58 children with a pediatric inflammatory multisystem syndrome temporally associated with SARS-CoV-2. *JAMA* 2020;324:259.
- 74 Lu X, Zhang L, Du H. SARS-CoV-2 infection in children. *N Engl J Med* 2020;382.
- 75 Göttinger F, Santiago-García B, Noguera-Julían A, *et al.* COVID-19 in children and adolescents in Europe: a multinational, multicentre cohort study. *Lancet Child Adolesc Health* 2020;4:653–61.
- 76 Parri N, Lenge M, Buonsenso D. Children with Covid-19 in pediatric emergency departments in Italy. *N Engl J Med Overseas Ed* 2020;383:187–90.
- 77 Abdel-Mannan O, Eyre M, Löbel U, *et al.* Neurologic and radiographic findings associated with COVID-19 infection in children. *JAMA Neurol* 2020. doi:10.1001/jamaneurol.2020.2687. [Epub ahead of print: 01 Jul 2020].
- 78 Banerjee S, Guha A, Das A, *et al.* A preliminary report of COVID-19 in children in India. *Indian Pediatr* 2020;57:963–4.
- 79 Bhumbra S, Malin S, Kirkpatrick L, *et al.* Clinical features of critical coronavirus disease 2019 in children. *Pediatr Crit Care Med* 2020;21:e948–e953.
- 80 Bisogno G, Provenzi M, Zama D. Clinical characteristics and outcome of SARS-CoV-2 infection in Italian pediatric oncology patients: a study from the infectious diseases Working group of the AIEOP. *J Pediatric Infect Dis Soc* 2020.
- 81 Blumfield E, Levin TL. COVID-19 in pediatric patients: a case series from the Bronx, NY. *Pediatr Radiol* 2020;50:1369–74.
- 82 Derespina KR, Kaushik S, Plichta A, *et al.* Clinical manifestations and outcomes of critically ill children and adolescents with coronavirus disease 2019 in New York City. *J Pediatr* 2020;226:10.1016/j.jpeds.2020.07.039. [Epub ahead of print: 16 Jul 2020].
- 83 George S, Ansari MS, Kalliath A. COVID-19 in children in Brunei Darussalam: higher incidence but mild manifestations. *J Med Virol* 2020.
- 84 González Cortés R, García-Salido A, Roca Pascual D, *et al.* A multicenter national survey of children with SARS-CoV-2 infection admitted to Spanish pediatric intensive care units. *Intensive Care Med* 2020;46:1774–6.
- 85 Kainth MK, Goenka PK, Williamson KA, *et al.* Early experience of COVID-19 in a US children's Hospital. *Pediatrics* 2020;146:e202003186.
- 86 Kaushik S, Aydin SI, Derespina KR, *et al.* Multisystem inflammatory syndrome in children associated with severe acute respiratory syndrome coronavirus 2 infection (MIS-C): a multi-institutional study from New York City. *J Pediatr* 2020;224:24–9.
- 87 Kest H, Kaushik A, DeBruin W. Multisystem inflammatory syndrome in children (MIS-C) associated with 2019 novel coronavirus (SARS-CoV-2) infection. *Case Rep Pediatr* 2020;2020:1–4.
- 88 Korkmaz MF, Türe E, Dorum BA, *et al.* The epidemiological and clinical characteristics of 81 children with COVID-19 in a pandemic hospital in turkey: an observational cohort study. *J Korean Med Sci* 2020;35:e236.
- 89 Meslin P, Guiomard C, Chouakria M, *et al.* Coronavirus disease 2019 in newborns and very young infants: a series of six patients in France. *Pediatr Infect Dis J* 2020;39:e145–7.
- 90 Moraleda C, Serna-Pascual M, Soriano-Arandes A, *et al.* Multi-Inflammatory syndrome in children related to SARS-CoV-2 in Spain. *Clin Infect Dis* 2020. doi:10.1093/cid/ciaa1042. [Epub ahead of print: 25 Jul 2020].
- 91 KF N, Kothari T, Bandi S. COVID-19 multisystem inflammatory syndrome in three teenagers with confirmed SARS-CoV-2 infection. *J Med Virol* 2020;92.
- 92 Paquette D, Bell C, Roy M, *et al.* Laboratory-confirmed COVID-19 in children and youth in Canada, January 15-April 7, 2020. *Can Commun Dis Rep* 2020;46:121–4.
- 93 Perez A, Kogan-Liberman D, Sheflin-Findling S. Presentation of severe acute respiratory syndrome-coronavirus 2 infection as cholestatic jaundice in two healthy adolescents. *J Pediatr* 2020. doi:10.1016/j.jpeds.2020.07.054. [Epub ahead of print: 23 Jul 2020].
- 94 Sarangi B, Reddy VS, Oswal JS, *et al.* Epidemiological and clinical characteristics of COVID-19 in Indian children in the initial phase of the pandemic. *Indian Pediatr* 2020;57:914–7.
- 95 Sadiq M, Aziz OA, Kazmi U, *et al.* Multisystem inflammatory syndrome associated with COVID-19 in children in Pakistan. *Lancet Child Adolesc Health* 2020;4:e36–7.
- 96 Swann OV, Holden KA, Turtle L, *et al.* Clinical characteristics of children and young people admitted to hospital with covid-19 in United Kingdom: prospective multicentre observational cohort study. *BMJ* 2020;370:m3249.
- 97 Antúnez-Montes OY, Escamilla MI, Figueroa-Urbe AF, *et al.* COVID-19 and multisystem inflammatory syndrome in Latin American children. *Pediatr Infect Dis J* 2020; Publish Ahead of Print.
- 98 Capone CA, Subramony A, Sweberg T, *et al.* Characteristics, cardiac involvement, and outcomes of multisystem inflammatory syndrome of childhood associated with severe acute respiratory syndrome coronavirus 2 infection. *J Pediatr* 2020;224:141–5.
- 99 Cura Yayla BC, Özsüreççi Y, Aykaç K, *et al.* Characteristics and management of children with COVID-19 in turkey. *Balkan Med J* 2020;37:341–7.
- 100 Davies P, Evans C, Kanthimathinathan HK, *et al.* Intensive care admissions of children with paediatric inflammatory multisystem syndrome temporally associated with SARS-CoV-2 (PIMS-TS) in the UK: a multicentre observational study. *Lancet Child Adolesc Health* 2020;4:669–77.



- 101 de Farias ECF, Pedro Piva J, de Mello MLFMF, *et al.* Multisystem inflammatory syndrome associated with coronavirus disease in children: a Multi-centered study in Belém, Pará, Brazil. *Pediatr Infect Dis J* 2020;39:e374-e376.
- 102 Dima M, Enatescu I, Craina M, *et al.* First neonates with severe acute respiratory syndrome coronavirus 2 infection in Romania: three case reports. *Medicine* 2020;99:e21284.
- 103 Klompas M, Baker MA, Rhee C. Airborne transmission of SARS-CoV-2: theoretical considerations and available evidence. *JAMA* 2020.
- 104 Ibrahim LF, Tosif S, McNab S, *et al.* SARS-CoV-2 testing and outcomes in the first 30 days after the first case of COVID-19 at an Australian children's hospital. *Emerg Med Australas* 2020;32:801-8.
- 105 Kilani MM, Odeh MM, Shalabi M, *et al.* Clinical and laboratory characteristics of SARS-CoV-2-infected paediatric patients in Jordan: serial RT-PCR testing until discharge. *Paediatr Int Child Health* 2020:1-10.
- 106 Okarska-Napierała M, Ludwikowska KM, Szenborn L, *et al.* Pediatric inflammatory multisystem syndrome (PIMS) did occur in Poland during months with low COVID-19 prevalence, preliminary results of a nationwide register. *J Clin Med* 2020;9. doi:10.3390/jcm9113386. [Epub ahead of print: 22 Oct 2020].
- 107 Mahmoudi S, Mehdi-zadeh M, Shervin Badv R, *et al.* The coronavirus disease 2019 (COVID-19) in children: a study in an Iranian children's referral hospital. *Infect Drug Resist* 2020;13:2649-55.
- 108 Mamishi S, Movahedi Z, Mohammadi M, *et al.* Multisystem inflammatory syndrome associated with SARS-CoV-2 infection in 45 children: a first report from Iran. *Epidemiol Infect* 2020;148:e196.
- 109 Mithal LB, Machut KZ, Muller WJ, *et al.* SARS-CoV-2 Infection in Infants Less than 90 Days Old. *J Pediatr* 2020;224:150-2.
- 110 Newman AM, Jhaveri R, Patel AB, *et al.* Trisomy 21 and coronavirus disease 2019 in pediatric patients. *J Pediatr* 2021;228:294-6.
- 111 Torres JP, Izquierdo G, Acuña M, *et al.* Multisystem inflammatory syndrome in children (MIS-C): report of the clinical and epidemiological characteristics of cases in Santiago de Chile during the SARS-CoV-2 pandemic. *Int J Infect Dis* 2020;100:75-81.
- 112 Saha S, Ahmed ANU, Sarkar PK, *et al.* The direct and indirect impact of SARS-CoV-2 infections on neonates: a series of 26 cases in Bangladesh. *Pediatr Infect Dis J* 2020;39:e398-e405.
- 113 Gale C, Quigley MA, Placzek A, *et al.* Characteristics and outcomes of neonatal SARS-CoV-2 infection in the UK: a prospective national cohort study using active surveillance. *Lancet Child Adolesc Health* 2021;5:113-21.
- 114 Schwartz DA, Mohagheghi P, Beigi B. Spectrum of neonatal COVID-19 in Iran: 19 infants with SARS-CoV-2 perinatal infections with varying test results, clinical findings and outcomes. *The Journal of Maternal-Fetal & Neonatal Medicine* 2020;2:1-10.
- 115 Schwartz SP, Thompson P, Smith M, *et al.* Convalescent plasma therapy in four critically ill pediatric patients with coronavirus disease 2019: a case series. *Crit Care Explor* 2020;2:e0237.
- 116 Almoosa ZA, Al Ameer HH, AlKadhem SM, *et al.* Multisystem Inflammatory Syndrome in Children, the Real Disease of COVID-19 in Pediatrics - A Multicenter Case Series From Al-Ahsa, Saudi Arabia. *Cureus* 2020;12:e11064.
- 117 Alsharrah DY, Al-Haddad F, Aljamaan S, *et al.* 441. clinical characteristics of pediatric SARS-CoV-2 infection and coronavirus disease 2019 (COVID-19) in Kuwait. *Open Forum Infect Dis* 2020;7:S288.
- 118 Prata-Barbosa A, Lima-Setta F, Santos GRD, *et al.* Pediatric patients with COVID-19 admitted to intensive care units in Brazil: a prospective multicenter study. *J Pediatr* 2020;96:582-92.
- 119 Bayesheva D, Boranbayeva R, Turdalina B, *et al.* COVID-19 in the paediatric population of Kazakhstan. *Paediatr Int Child Health* 2020:1-7.
- 120 Bhavsar SM, Clouser KN, Gadhavi J, *et al.* COVID-19 in pediatrics: characteristics of hospitalized children in New Jersey. *Hosp Pediatr* 2021;11:79-87.
- 121 Bustos-Cordova E, Castillo-García D, Cerón-Rodríguez M, *et al.* Clinical spectrum of COVID-19 in a Mexican pediatric population. *Indian Pediatr* 2020. [Epub ahead of print: 19 Dec 2020].
- 122 Fernández Colomer B, Sánchez-Luna M, de Alba Romero C, *et al.* Neonatal infection due to SARS-CoV-2: an epidemiological study in Spain. *Frontiers in Pediatrics* 2020;8:670.
- 123 Storch-de-Gracia P, Leoz-Gordillo I, Andina D, *et al.* Clinical spectrum and risk factors for complicated disease course in children admitted with SARS-CoV-2 infection. *Anales de Pediatría* 2020;93:323-33.
- 124 Dhanalakshmi K, Venkataraman A, Balasubramanian S, *et al.* Epidemiological and Clinical Profile of Pediatric Inflammatory Multisystem Syndrome - Temporally Associated with SARS-CoV-2 (PIMS-TS) in Indian Children. *Indian Pediatr* 2020;57:1010-4.
- 125 Falah NU, Hashmi S, Ahmed Z, *et al.* Kawasaki disease-like features in 10 pediatric COVID-19 cases: a retrospective study. *Cureus* 2020;12:e11035.
- 126 García-Salido A, de Carlos Vicente JC, Belda Hofheinz S, *et al.* Severe manifestations of SARS-CoV-2 in children and adolescents: from COVID-19 pneumonia to multisystem inflammatory syndrome: a multicentre study in pediatric intensive care units in Spain. *Crit Care* 2020;24:1-13.
- 127 Jain S, Sen S, Lakshmvikenateshiah S, *et al.* Multisystem inflammatory syndrome in children with COVID-19 in Mumbai, India. *Indian Pediatr* 2020;57:1015-9.
- 128 Kalamdani P, Kalathingal T, Manerker S, *et al.* Clinical profile of SARS-CoV-2 infected neonates from a tertiary government hospital in Mumbai, India. *Indian Pediatr* 2020;57:1143-6.
- 129 Lima-Setta F, Magalhães-Barbosa MCde, Rodrigues-Santos G, *et al.* Multisystem inflammatory syndrome in children (MIS-C) during SARS-CoV-2 pandemic in Brazil: a multicenter, prospective cohort study. *J Pediatr* 2020. doi:10.1016/j.jpeds.2020.10.008. [Epub ahead of print: 09 Nov 2020].
- 130 Montoya J, Ugaz C, Alarcon S, *et al.* COVID-19 in pediatric cancer patients in a resource-limited setting: national data from Peru. *Pediatr Blood Cancer* 2021;68:e28610.
- 131 Önal P, Kılınç AA, Ayygün F. COVID-19 in turkey: a tertiary center experience. *Pediatr Int* 2020.
- 132 Rabha AC, Oliveira Junior Fde, Oliveira TAdE, *et al.* Clinical manifestations of children and adolescents with covid-19: report of the first 115 cases from sabará hospital infantil. *Revista Paulista de Pediatria* 2021;39.
- 133 Sethuraman U, Kannikeswaran N, Ang J, *et al.* Multisystem inflammatory syndrome in children associated with novel coronavirus SARS-CoV-2: presentations to a pediatric emergency department in Michigan. *Am J Emerg Med* 2021;39:164-7.
- 134 Shahbaznejad L, Navaeifar MR, Abbaskhanian A, *et al.* Clinical characteristics of 10 children with a pediatric inflammatory multisystem syndrome associated with COVID-19 in Iran. *BMC Pediatr* 2020;20:1-12.
- 135 Webb K, Abraham DR, Faley A, *et al.* Multisystem inflammatory syndrome in children in South Africa. *Lancet Child Adolesc Health* 2020;4:e38.
- 136 Zachariah P, Johnson CL, Halabi KC, *et al.* Epidemiology, clinical features, and disease severity in patients with coronavirus disease 2019 (COVID-19) in a children's hospital in New York City, New York. *JAMA Pediatr* 2020;174:e202430-e.
- 137 van der Zalm MM, Lishman J, Verhagen LM. *Clinical experience with SARS CoV-2 related illness in children-hospital experience in Cape town, South Africa. Clinical infectious diseases: an official publication of the infectious diseases Society of America*, 2020.
- 138 Zheng G, Wang B, Zhang H, *et al.* Clinical characteristics of acute respiratory syndrome with SARS-CoV-2 infection in children in South China. *Pediatr Pulmonol* 2020;55:2419-26.
- 139 Jiang L, Tang K, Levin M, *et al.* COVID-19 and multisystem inflammatory syndrome in children and adolescents. *Lancet Infect Dis* 2020;20:e276-88.
- 140 Ludvigsson JF. Systematic review of COVID-19 in children shows milder cases and a better prognosis than adults. *Acta Paediatr* 2020;109:1088-95.
- 141 Huang C, Wang Y, Li X, *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395:497-506.
- 142 Terry C, Mühlemann B, Veith T, *et al.* An analysis of SARS-CoV-2 viral load by patient age 2020.
- 143 Chang T-H, Wu J-L, Chang L-Y. Clinical characteristics and diagnostic challenges of pediatric COVID-19: a systematic review and meta-analysis. *J Formos Med Assoc* 2020;119:982-989.
- 144 NBS. *National Bureau of statistics of China*, 2019.
- 145 Ludvigsson JF. Case report and systematic review suggest that children may experience similar long-term effects to adults after clinical COVID-19. *Acta Paediatr* 2020;370.

**Clinical characteristics, treatment and outcomes of pediatric COVID-19: a systematic review and meta-analysis****Supplement file**

**Table 1: Search Strategy for MEDLINE**

**Table 2: Distribution of studies (N=129) according to Country of origin included in the review**

**Table 3: Characteristics of included studies (N=129)**

**Table 4: Age Distribution from 91 studies of disease prevalence**

**Table 5: Age Distribution from 67 case series of clinical features and outcomes**

**Table 6: Characteristics of included studies with MIS-C**

**Table 7: Characteristics of included studies from neonatal age group**

**Table 8: Quality assessment of included studies (N=129)**

**Figure 1: Flow diagram of the systematic review process**

**Figure 2: Global map of distribution of the studies included (N=129) in the review**

**Table 1: Search Strategy for MEDLINE**

- 1- (coronavirus or "covid 2019" or "SARS2" or "SARS" or "SARS CoV19" or "severe acute respiratory syndrome coronavirus 2" or "coronavirus infection" or "severe acute respiratory pneumonia outbreak" or "novel cov" or 2019ncov or sars cov2 or cov2 or ncov or covid or covid19 or coronaviridae or "corona virus").mp. [mp=title, abstract, original title, name of substance word, subject heading word, floating sub-heading word, keyword heading word, organism supplementary concept word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
- 2- "coronavirus infections".mp. or exp Coronavirus Infections/
- 3- exp Coronavirus 229E, Human/ or exp Coronavirus/ or exp Middle East Respiratory Syndrome Coronavirus/ or coronavirus.mp.
- 4- 1 or 2 or 3
- 5- (infant or neonate or newborn or child or children or pediatric\* or paediatric or baby or babies or toddler).mp. [mp=title, abstract, original title, name of substance word, subject heading word, floating sub-heading word, keyword heading word, organism supplementary concept word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
- 6- exp Infant, Low Birth Weight/ or exp Infant, Extremely Premature/ or exp Infant/ or exp Infant, Newborn/ or exp Infant, Premature/
- 7- 5 or 6
- 8- 4 and 7

**Table 2: Distribution of studies (N=129) according to Country of origin included in the review**

Country	Count of reports	Sum of COVID-19 cases
USA	24	3443
China	36	1315
Canada	1	938
UK	9	867
Kazakhstan	1	650
25 European Countries	1	582
Italy	4	171
Spain	8	287
Korea	1	91
India	5	145
Turkey	3	338
Jordan	1	61
France	4	47
Iran	7	152
Brunei-Darussalam	1	12
Chile	1	27
Bangladesh	1	26
Brazil	4	261
Pakistan	3	20
Poland	1	9
Kuwait	1	134
Saudi Arabia	1	10
Australia	1	4
Malaysia	1	4
Romania	1	3
Japan	1	3
Germany	1	3
Mexico	2	53
South Africa	2	182
Peru	1	69
5 Latin American Countries (Mexico, Colombia, Peru, Costa Rico, Brazil)	1	409

Total

129

10251

Table 3: Characteristics of included studies (N=129)

Study and year	Country	World Bank Country Classification	Total number (N)	Study setting	Mean age (years)	Males (N)	Past medical history and Contact history	Presenting signs and symptoms	Management (other than supportive care)	Outcomes
Qiu, H 2020	China	LMIC	36	Patients presenting to hospital	8.3	23	Travel history (n=12) Contact history (n=32) Comorbidities (NR)	Fever (n=13), cough (n=7), headache (n=3), nasal symptoms (n=2), GI symptoms (n=2)	Interferon (n=36), antivirals (n=14), oxygen (n=6)	Recovered (n=36) ICU admission (n=0)
Wang, D 2020	China	LMIC	31	Patients admitted in hospital	7.1	15	Travel history (n=9) Contact history (n=1) Comorbidities (n=0)	Asymptomatic (n=7), fever (n=20), cough (n=14), fatigue and diarrhea (n=3 each), sore throat (n=2), headache/dizziness (n=3), rhinorrhoea (n=2), vomiting (n=2)	Interferon (n=10), Antibiotics (n=6), antivirals (n=29), traditional medicine (n=9), IVIG (n=4)	Recovered (n=24) ICU admission (n=0)
Zheng, F 2020	China	LMIC	25	Patients admitted in hospital	3.0	14	Travel history (n=9) Contact history (n=1) Comorbidities (n=0)	Fever (n=13), cough (n=11), diarrhea (n=3), dyspnea (n=2), vomiting (n=2), abdominal pain (n=2), nasal congestion (n=2)	Interferon (n=2), Antibiotics (n=2), antivirals (n=2), steroids (n=2), IVIG (n=4), Mechanical ventilation (n=2)	Recovered (NR) ICU admission (n=2)

Xia, W 2020	China	LMIC	20	Patients admitted in hospital	4.8	13	Travel history (n=20) Contact history (n=13) Comorbidities (NR)	Cough (n=13), fever (n=12), diarrhea (n=3), dyspnea (n=2), sore throat (n=1), fatigue (n=1)	NR	Recovered (n=19) ICU admission (n=1)
Feng, K 2020	China	LMIC	15	Patients presenting to hospital	NR	5	Travel history (n=3) Contact history (n=12) Comorbidities (NR)	Asymptomatic (n=8), fever (n=5), cough or nasal congestion (n=1)	NR	Recovered (n=5) ICU admission (n=0)
Cai, J 2020	China	LMIC	10	Patients admitted in hospital	6.5	4	NR	Fever (n=8), cough (n=6) or sore throat or nasal congestion (n=7)	Antibiotics (n=5)	Recovered (n=10) ICU admission (n=0)
Wei, M 2020	China	LMIC	9	Patients admitted in hospital	0.5	2	Travel history (n=7) Contact history (n=9) Comorbidities (NR)	Asymptomatic (n=6), fever (n=4), cough (n=2), rhinorrhoea (n=1), sputum production (n=1)	NR	Recovered (n=9) ICU admission (n=0)
Su, L 2020	China	LMIC	9	Patients presenting to hospital	4.5	3	Travel history (n=8) Contact history (n=9) Comorbidities (NR)	Asymptomatic (n=6), fever or cough (n=3)	Interferon given to all patients; antivirals (n=2)	Recovered (NR) ICU admission (n=0)
Zhou, Y 2020	China	LMIC	9	Patients admitted in hospital	1.0	4	Travel history (n=3) Contact history (n=9) Comorbidities (n=0)	Asymptomatic (n=5), fever (n=4), cough (n=2), rhinorrhoea (n=1)	Interferon (n=9) antivirals (n=6), mechanical ventilation (n=9)	Recovered (NR) ICU admission (n=0)
Sun, D 2020	China		8		6.8	6	Travel history (n=8)	Fever or cough (n=6 each), sputum production (n=4),	mechanical ventilation (n=2), antibiotics (n=5),	Recovered (n=5)

		LMIC		Patients admitted in hospital			Contact history (n=5) Comorbidities (n=3)	nausea/ vomiting (n=4), diarrhea (n=3), fatigue (n=1) or headache (n=1), sore throat or nasal symptoms (n=8)	systemic steroids (n=5), IVIG (n=4), Chinese medications (n=4)	ICU admission (n=8)
Liu, W 2020	China	LMIC	6	Patients admitted in hospital	3.0	2	NR	High fever and cough in all patients; vomiting and fatigue (n=4)	Antivirals (n=6), systemic steroids (n=4), IVIG (n=1)	Recovered (n=5) ICU admission (n=1)
Hu, Z 2020	China	LMIC	7	Patients admitted in hospital	8.0	5	Travel history (n=3) Contact history (n=0) Comorbidities (NR)	Asymptomatic (n=4), fever (n=1)	NR	Recovered (n=6) ICU admission (NR)
Liu, M 2020	China	LMIC	5	Patients admitted in hospital	5.9	4	Travel history (n=0) Contact history (n=5) Comorbidities (NR)	Fever (n=2), cough (n=2), diarrhea (n=1), asymptomatic (n=3)	Antiviral (n=3), Interferon (n=4)	Recovered (n=5) ICU admission (n=0)
Liu, H 2020	China	LMIC	4	Patients presenting to hospital	3.0	2	Travel history (n=4) Contact history (n=4) Comorbidities (NR)	Fever (n=3), cough (n=3), Fatigue (n=1)	NR	NR
Lou, MX 2020	China	LMIC	3	Patients admitted in hospital	4.8	1	Travel history (n=2) Contact history (n=3) Comorbidities (NR)	Fever (n=3), cough (n=1), Fatigue (n=2), diarrhea (n=2), nasal symptoms or sore throat (n=2)	Antibiotics (n=1), interferon (n=2)	Recovered (n=3) ICU admission (n=0)
Zhang, T 2020	China	LMIC	3	Patients admitted in hospital	7.7	3	Travel history (n=0) Contact history (n=3)	Fever (n=2), cough (n=1), Fatigue (n=1), nausea or vomiting (n=2), nasal symptoms or sore throat (n=3)	Antibiotics (n=1), antivirals (n=1), interferon (n=3), Chinese medicine (n=3)	Recovered (n=3) ICU admission (NR)

							Comorbidities (NR)			
Li, Y 2020	China	LMIC	2	Patients admitted in hospital	4.0	1	Travel history (n=2) Contact history (n=2) Comorbidities (n=0)	Cough (n=1), nasal symptoms (n=1), asymptomatic (n=1)	NR	Recovered (NR) ICU admission (NR)
Ji, LN 2020	China	LMIC	2	Patients admitted in hospital	12.0	2	Travel history (n=2) Contact history (n=2) Comorbidities (n=0)	Fever (n=1), nasal symptoms (n=1), sore throat (n=1)	NR	Recovered (NR) ICU admission (n=0)
Tagarro, A 2020	Spain	HIC	41	Patients admitted in hospital	4.3	18	Travel history (NR) Contact history (n=16) Comorbidities (n=2)	Upper respiratory symptoms in 14 (34%), fever (n=11), gastroenteritis or vomiting (n=2)	Mechanical ventilation (n=4)	Recovered (n=37) ICU admission (n=4)
Chen, A 2020	China	LMIC	14	Patients admitted in hospital	4.3	8	Travel history (n=10) Contact history (n=4) Comorbidities (n=0)	Fever (n=6), cough (n=6), Fatigue (n=3), nasal symptoms or sore throat (n=2)	NR	NR
See, KC 2020	Malaysia	LMIC	4	Patients presenting to hospital	6.4	3	Travel history (n=4) Contact history (n=0) Comorbidities (N=1)	Fever (n=1), cough (n=2), diarrhea (n=1), asymptomatic (n=2)	Antibiotics (n=1)	Recovered (n=4) ICU admission (n=0)
Zhang, GX 2020	China	LMIC	2	Patients admitted in hospital	1.2	0	Travel history (n=0) Contact history (n=2)	Fever (n=2), fatigue (n=1), dyspnea (n=1), diarrhea (n=2), nausea or vomiting (n=2),	IVIg (n=1), interferon (n=2), Chinese medicine (n=1)	Recovered (n=2) ICU admission (n=0)



							Comorbidities (n=0)			
Cheng, X 2020	China	LMIC	2	Patients admitted in hospital	1.0	1	Travel history (n=2) Contact history (n=0) Comorbidities (n=0)	Fever (n=2), fatigue (n=1), dyspnea (n=1), diarrhea (n=2), nausea or vomiting (n=2),	Antibiotics (n=2), interferon (n=1), Chinese medicine (n=1)	Recovered (n=2) ICU admission (n=1)
Tang, A 2020	China	LMIC	26	Patients admitted in hospital	6.9	17	Travel history (n=26) Contact history (n=26) Comorbidities (n=0)	Fever (n=11), cough (n=12), diarrhea (n=2), nausea or vomiting (n=2), nasal symptoms or sore throat (n=2), asymptomatic (n=9)	Antivirals (n=22), interferon (n=4)	Recovered (n=17) ICU admission (n=0)
Zhu, L 2020	China	LMIC	10	Patients admitted in hospital	9.1	1	Travel history (n=8) Contact history (n=10) Comorbidities (n=0)	Fever (n=3), cough (n=3), asymptomatic (n=6)	Antivirals (n=1), interferon (n=10)	Recovered (n=10) ICU admission (n=0)
Tan, X 2020	China	LMIC	13	Patients admitted in hospital	7.9	4	Travel history (n=3) Contact history (n=10) Comorbidities (n=0)	Fever (n=6), cough (n=6), nasal symptoms (n=1), diarrhea (n=1), nausea or vomiting (n=1), abdominal pain (n=1), headache (n=1), asymptomatic (n=2)	Antivirals (n=12), interferon (n=10)	Recovered (n=13) ICU admission (n=0)
Ma, H 2020	China	LMIC	22	Patients admitted in hospital	5.5	12	Travel history (n=22) Contact history (n=0) Comorbidities (n=0)	Fever (n=13), dry cough (n=5), shortness of breath (n=1), asymptomatic (n=2)	NR	Recovered (n=5) ICU admission (n=0)
Shen, Q 2020	China	LMIC	9	Patients admitted in hospital	7.6	3	Travel history (n=2) Contact history (n=6)	asymptomatic (n=2), fever (n=3), diarrhea (n=2), sore throat (n=1), cough (n=1)	Antivirals (n=9), antibiotics (n=4), IVIG (n=1), Systemic steroids (n=1)	Recovered (n=6) ICU admission (n=0)

							Comorbidities (n=0)			
Schwierz eck, V 2020	Germany	HIC	3	Patients admitted in hospital	10.0	NR	Travel history (n=0) Contact history (NR) Comorbidities (n=3)	Asymptomatic (n=2), fever (n=1), cough (n=1), nasal congestion (n=1)	NR	NR
Jiang, JY 2020	China	LMIC	6	Patients presenting to hospital	7.0	5	Travel history (n=0) Contact history (n=6) Comorbidities (NR)	Fever (n=3), cough (n=2), diarrhea (n=3), asymptomatic (n=3)	NR	Recovered (n=6) ICU admission (n=0)
Riphagen , S 2020	UK	HIC	8	Patients admitted in hospital	8.9	5	Travel history (NR) Contact history (NR) Comorbidities (NR)	Fever (n=8), diarrhea (n=7), rashes (n=4), conjunctivitis (n=5)	IVIg (n=6), Systemic steroids (n=5), Inotropes (n=8), aspirin (n=3)	Recovered (n=7) ICU admission (n=8) Death (n=1)
Shekerde mian, LS 2020	USA	HIC	48	Patients admitted to intensive care units	11.7	25	Travel history (NR) Contact history (NR) Comorbidities (n=40)	Multi-organ failure (n=11), pulmonary symptoms (n=35), diarrhea (n=1), headache (n=2), asymptomatic (n=1)	hydroxychloroquine (n=21), antiviral (n=8), mechanical ventilation (n=18), antibiotics (n=15)	Recovered (n=31) ICU admission (n=48) Death (n=2)
Verdoni, L 2020	Italy	HIC	10	Patients admitted in hospital	7.5	7	Travel history (NR) Contact history (NR) Comorbidities (n=5)	GI symptoms (n=5), neurological (n=4), Kawasaki signs (n=12), rashes (n=5)	IVIg (n=8), Inotropes (n=2)	Recovered (n=10) ICU admission (n=0)
Rahimza deh, G 2020	Iran	LMIC	9	Patients admitted in hospital	4.9	6	Travel history (n=0) Contact history (n=0)	Fever (n=9), cough (n=9), fatigue (n=9), asymptomatic (n=3)	Mechanical ventilation (n=2), antivirals (n=9), antibiotics (n=9), antimalarials (n=9)	Recovered (n=9) ICU admission (n=2)

							Comorbidities (n=0)			
Tullie, L 2020	UK	HIC	8	Patients admitted in hospital	NR	NR	NR	Fever (n=8), Kawasaki signs (n=4), rashes (n=4), GI symptoms (n=8)	antibiotics (n=8), IVIG (n=4)	Recovered (n=7) ICU admission (n=4)
Sajid, MI 2020	Pakistan	LMIC	2	Patients admitted in hospital	12.0	1	Travel history (NR) Contact history (NR) Comorbidities (n=2)	Fever (n=2), cough (n=1)	NR	Recovered (NR) ICU admission (n=0)
Zhang, ZJ 2020	China	LMIC	4	Patients admitted in hospital	Newborns	3	Travel history (NR) Contact history (NR) Comorbidities (n=2)	Fever (n=2), shortness of breath (n=1), cough (n=1), vomiting (n=1), asymptomatic (n=1)	NR	Recovered (NR) ICU admission (n=0)
Zeng, L 2020	China	LMIC	3	Patients admitted in hospital	Newborns	3	Travel history (n=3) Contact history (n=3) Comorbidities (n=0)	Fever (n=2), cough (n=1), fatigue (n=1)	Mechanical ventilation (n=1), antibiotics (n=1)	Recovered (NR) ICU admission (n=3)
Salvatori, G 2020	Italy	HIC	2	Patients presenting to hospital	Newborns	1	Travel history (NR) Contact history (NR) Comorbidities (n=0)	Cough (n=1), fatigue (n=1), diarrhea (n=1), asymptomatic (n=1)	NR	Recovered (n=2) ICU admission (n=0)
Zhang, X 2020	China	LMIC	10	Patients presenting to hospital	4.9	3	Travel history (n=0) Contact history (n=10) Comorbidities (n=0)	Fever (n=3), Cough (n=1), sore throat (n=1), asymptomatic (n=5)	NR	Recovered (n=10) ICU admission (n=0)
Wu, HP 2020	China		23		NR	NR	Travel history (n=7)	Fever (n=16), Cough (n=11), diarrhea (n=2),	Antivirals (n=8), antibiotics (n=6),	Recovered (n=23) ICU admission (n=0)

		LMIC		Patients admitted in hospital			Contact history (n=17) Comorbidities (n=0)	sore throat (n=2), asymptomatic (n=3)	interferon (n=17), antimalarial (n=1)	
Zhang, H 2020	China	LMIC	2	Patients presenting to hospital	13.5	2	Travel history (n=2) Contact history (n=2) Comorbidities (n=2)	Fever (n=1), Cough (n=1), asymptomatic (n=1)	Antivirals (n=2), antibiotics (n=1), interferon (n=2)	Recovered (n=2) ICU admission (n=0)
Chao, JY 2020	USA	HIC	46	Patients admitted in hospital	13.1	31	Travel history (n=0) Contact history (n=20) Comorbidities (n=31)	Fever (n=28), Cough (n=28), dyspnea (n=22) asymptomatic (n=1)	NR	Recovered (n=45) ICU admission (n=13) Deaths (n=1)
Eghbali, A 2020	Iran	LMIC	4	Patients admitted in hospital	11.0	4	Travel history (NR) Contact history (NR) Comorbidities (n=2)	Fever (n=4), Cough (n=4), dyspnea (n=2)	Mechanical ventilation (n=2)	Recovered (n=4) ICU admission (n=2)
Chiotos, K 2020	USA	HIC	6	Patients admitted in hospital	8.5	1	Travel history (n=0) Contact history (n=0) Comorbidities (n=0)	Fever (n=6), diarrhea (n=6), nausea of vomiting (n=5), abdominal pain (n=5), dyspnea (n=4), kawasaki signs (n=6), rashes (n=2), conjunctivitis (n=2)	IVIG (N=6), Systemic steroids (n=3), inotropes (n=6), aspirin (n=3)	Recovered (n=5) ICU admission (n=1)
Flores, V 2020	Mexico	LMIC	3	Patients admitted in hospital	7.0	0	Travel history (NR) Contact history (n=0) Comorbidities (n=3)	Fever (n=3), dyspnea (n=1)	Mechanical ventilation (n=1), Antibiotics (n=3)	Recovered (n=2) ICU admission (n=1) Deaths (n=1)
García-Salido, A 2020	Spain		7	Patients admitted in hospital	8.4	4	Travel history (NR)	Fever (n=5), dyspnea (n=3), diarrhea (n=3), nausea or vomiting (n=3)	Antivirals (n=6), antibiotics (n=6), IVIG	Recovered (n=7) ICU admission (n=7)

		HIC		Patients admitted to intensive care units			Contact history (NR) Comorbidities (n=1)		(n=2), systemic steroids (n=6), antimalarial (n=6), interferon (n=2)	
Cabrero-Hernández, M 2020	Spain	HIC	5	Patients admitted to intensive care units	10.6	4	Travel history (NR) Contact history (NR) Comorbidities (n=0)	Fever (n=5), diarrhea (n=3), nausea or vomiting (n=2), abdominal pain (n=5), Kawasaki signs (n=1), rashes (n=1)	Mechanical ventilation (n=5), Systemic steroids (n=5), antimalarial (n=5), inotropes (n=4)	Recovered (n=5) ICU admission (n=5)
Waltuch, T 2020	USA	HIC	4	Patients admitted to intensive care units	10.0	3	Travel history (n=0) Contact history (n=2) Comorbidities (n=2)	Fever (n=4), Cough (n=2), fatigue (n=2), GI symptoms (n=3), Kawasaki signs (n=2), rashes (n=2), conjunctivitis (n=1)	Antibiotics (n=4), IVIG (n=2), Inotropes (n=1)	Recovered (n=4) ICU admission (n=4)
Kakuya, F 2020	Japan	HIC	3	Patients presenting to hospital	8.3	3	Travel history (n=0) Contact history (n=1) Comorbidities (n=0)	Fever (n=3), Cough (n=3), fatigue (n=1),	Antibiotics (n=2)	Recovered (n=3) ICU admission (n=0)
Soltani, J 2020	Iran	LMIC	30	Patients presenting to hospital	6.0	14	Travel history (NR) Contact history (NR) Comorbidities (n=11)	Fever (n=23), Cough (n=16), dyspnea (n=20), diarrhea (n=3), nausea or vomiting (n=8), nasal symptoms or sore throat (n=6)	Antivirals (n=19), antibiotics (n=17), IVIG (n=2), antimalarial (n=23)	Recovered (n=29) ICU admission (n=1) Deaths (n=1)
Toubiana, J 2020	France	HIC	21	Patients admitted in hospital	7.9	9	Travel history (NR) Contact history (NR) Comorbidities (n=0)	Fever (n=11), GI Symptoms (n=21), neurological (n=6), kawasaki signs (n=11), rashes (n=16), conjunctivitis (n=17)	Mechanical ventilation (n=11), Antibiotics (n=18), IVIG (N=21), Systemic steroids (n=10), Inotropes (n=15), aspirin (n=21)	Recovered (n=21) ICU admission (n=17)
Foster, CE 2020	USA	HIC	57	Patients presenting to hospital	10.6	31	Travel history (n=3) Contact history (n=30)	Fever (n=35), cough (n=34), GI Symptoms (n=21), sore throat or	Antibiotics (n=3), systemic steroids (n=3), IVIG (n=1),	Recovered (n=57) ICU admission (n=0)

							Comorbidities (n=21)	nasal symptoms (n=16), asymptomatic (n=3)		
White, A 2020	USA	HIC	3	Patients admitted in hospital	0.1	2	Travel history (n=0) Contact history (n=2) Comorbidities (n=0)	Fever (n=3), dyspnea (n=3), nasal symptoms (n=3)	NR	Recovered (n=3) ICU admission (n=3)
Lin, EE 2020	USA	HIC	12	Patients admitted in hospital	11.3	8	Travel history (n=0) Contact history (n=2) Comorbidities (n=0)	Fever (n=3), cough (n=1), GI Symptoms (n=1), sore throat (n=2),	NR	Recovered (NR) ICU admission (n=0)
Gao, Y 2020	China	LMIC	11	Patients presenting to hospital	10.3	6	Travel history (n=0) Contact history (n=10) Comorbidities (n=0)	Fever (n=2), cough (n=2), dyspnea (n=1), asymptomatic (n=8)	Antiviral (n=4), interferon (n=7), Chinese medicine (n=5)	Recovered (n=11) ICU admission (n=0)
Harman, K 2020	UK	HIC	5	Patients admitted in hospital	7.1	2	Travel history (NR) Contact history (NR) Comorbidities (n=5)	Fever (n=3), asymptomatic (n=2)	Antiviral (n=1), antibiotics (n=5), antimalarial (n=1)	Recovered (n=4) ICU admission (n=2)
Blondiaux, E 2020	France	HIC	4	Patients admitted in hospital	9.0	1	Travel history (NR) Contact history (n=3) Comorbidities (n=0)	Fever (n=4), GI symptoms (n=4), rashes (n=4), conjunctivitis (n=3)	Mechanical ventilation (n=1), systemic steroids (n=3), inotropes (n=3), aspirin (n=3), IVIG (N=4)	Recovered (n=4) ICU admission (n=4)
Pouletty, M 2020	France	HIC	16	Patients admitted in hospital	10.0	8	Travel history (n=0) Contact history (n=12)	Fever (n=16), cough (n=2), dyspnea (n=2), GI symptoms (n=13), neurological (n=9), Kawasaki signs (n=7),	Mechanical ventilation (n=2), systemic steroids (n=4), antimalarial (n=1), aspirin (n=3), IVIG	Recovered (n=16) ICU admission (n=6)

							Comorbidities (n=6)	rashes (n=13), conjunctivitis (n=15)	(N=15), inotropes (n=6), aspirin (n=15)	
Whittaker, E 2020	UK	HIC	45	Patients admitted in hospital	10.0	19	Travel history (NR) Contact history (NR) Comorbidities (n=7)	Fever (n=45), dyspnea (n=9), GI symptoms (n=25), neurological (n=5), Kawasaki signs (n=25), rashes (n=21), conjunctivitis (n=20)	Mechanical ventilation (n=20), systemic steroids (n=33), IVIG (N=33), inotropes (n=23)	Recovered (n=44) ICU admission (n=20) Death (n=1)
Cheung, EW 2020	USA	HIC	17	Patients admitted in hospital	8.0	8	Travel history (n=0) Contact history (n=11) Comorbidities (n=3)	Fever (n=17), cough (n=7), dyspnea (n=7), neurological (n=8), Kawasaki signs (n=13), rashes (n=12), conjunctivitis (n=11)	Systemic steroids (n=12), IVIG (N=13), inotropes (n=10), aspirin (n=4)	Recovered (n=17) ICU admission (n=15)
Lu, X 2020	China	LMIC	171	National surveillance	6.7	104	Travel history (n=17) Contact history (n=131) Comorbidities (n=9)	Fever (n=71), cough (n=83), pharyngeal erythema (n=79), fatigue (n=13), diarrhea (n=15), nausea or vomiting (n=11), nasal symptoms or sore throat (n=22), asymptomatic (n=27)	NR	Recovered (n=149) ICU admission (n=3) Death (n=1)
CDC, COVID 2020	USA	HIC	2572	National surveillance	9.8	1408	Travel history (n=16) Contact history (n=168) Comorbidities (n=80)	Fever (n=163), cough (n=158), dyspnea (n=39), diarrhea (n=37), nausea or vomiting (n=31), abdominal pain (n=17), nasal symptoms or sore throat (n=71), neurological (n=81)	NR	Recovered (n=2569) ICU admission (n=15) Death (n=3)
Gotzinger, F 2020	Europe	HIC	582	Multinational cohort study	5.6	311	Travel history (n=0) Contact history (n=348) Comorbidities (n=145)	Fever (n=379), GI symptoms (n=128), neurological (n=70), asymptomatic (n=92)	Antiviral (n=30), IVIG (n=7), systemic steroids (n=22), antimalarial (n=40)	Recovered (n=460) ICU admission (n=48) Death (n=4)

Dong, Y 2020	China	LMIC	728	National surveillance study	9.8	418	Travel history (NR) Contact history (NR) Comorbidities (NR)	315 mild illness, 300 moderate illness, 18 severe and 3 critically ill, 94 asymptomatic children	NR	Recovered (n=727) ICU admission (n=21) Death (n=1)
Abdel-Mannan, O 2020	UK	HIC	4	Patients admitted in hospital	11.8	2	Travel history (NR) Contact history (NR) Comorbidities (NR)	Fever (n=4), dyspnea (n=2), nausea or vomiting (n=4), abdominal pain (n=1), neurological (n=4), rashes (n=4)	Mechanical ventilation (n=4), IVIG (n=3), systemic steroids (n=2)	Recovered (n=2) ICU admission (n=4)
Guha, A 2020	India	LMIC	41	National surveillance	1.9	24	Travel history (n=0) Contact history (n=26) Comorbidities (n=25)	Fever (n=9), cough (n=5), dyspnea (n=13), diarrhea (n=3), Kawasaki signs (n=2), rashes (n=2), asymptomatic (n=11)	Mechanical ventilation (n=2)	Recovered (n=40) ICU admission (n=6) Death (n=1)
Bhumbra, S 2020	USA	HIC	19	Patients admitted in hospital	6.7	14	Travel history (NR) Contact history (NR) Comorbidities (n=8)	Fever (n=12), cough (n=5), dyspnea (n=12), nausea or vomiting (n=2), sore throat (n=2),	Mechanical ventilation (n=4)	Recovered (n=12) ICU admission (n=7) Death (n=1)
Bisogno, G 2020	Italy	HIC	29	Patients admitted in hospital	7.5	13	Travel history (NR) Contact history (NR) Comorbidities (n=29)	Fever (n=8), fatigue (n=1), dyspnea (n=7), diarrhea (n=1), nausea or vomiting (n=1), asymptomatic (n=7)	Mechanical ventilation (n=0), antiviral (n=3), antibiotics (n=13), antimalarial (n=9)	Recovered (n=27) ICU admission (n=0)
Blumfield, E 2020	USA	HIC	19	Patients admitted in hospital	8.5	10	Travel history (NR) Contact history (NR)	Fever (n=17), cough (n=13), dyspnea (n=13), diarrhea (n=9), nausea or vomiting (n=9), abdominal pain (n=2),	Mechanical ventilation (n=8)	Recovered (n=17) ICU admission (n=14) Death (n=2)



							Comorbidities (n=12)	neurological (n=5), nasal symptoms or sore throat (n=8)		
Derespina, KR 2020	USA	HIC	70	Patients admitted in hospital	14.5	43	Travel history (n=2) Contact history (n=30) Comorbidities (n=52)	Fever (n=51), cough (n=50), dyspnea (n=45), fatigue (n=13), diarrhea (n=18), nausea or vomiting (n=24), neurological (n=15), nasal symptoms or sore throat (n=9)	Mechanical ventilation (n=20), antiviral (n=13), antibiotics (n=23), systemic steroids (n=23), antimalarial (n=27), inotropes (n=14)	Recovered (n=55) ICU admission (n=2) Death (n=70)
Feldstein, LR 2020	USA	HIC	186	Patients admitted in hospital	8.1	115	Travel history (NR) Contact history (NR) Comorbidities (n=51)	Fever (n=186), GI symptoms (n=170), neurological (n=10), Kawasaki signs (n=186), rashes (n=110), conjunctivitis (n=103)	Mechanical ventilation (n=37), IVIG (n=144), systemic steroids (n=91),	Recovered (n=130) ICU admission (n=148) Death (n=4)
George, S 2020	Brunei-Dar	LMIC	12	National surveillance	6.1	7	Travel history (n=0) Contact history (n=12) Comorbidities (n=4)	Fever (n=6), cough (n=6), asymptomatic (n=4)	Antibiotics (n=1)	Recovered (n=12) ICU admission (n=0)
Gonzalez-Cortes, R 2020	Spain	HIC	50	Patients admitted to intensive care units	6.7	31	Travel history (NR) Contact history (NR) Comorbidities (NR)	Dyspnea (n=20), neurological (n=1), Kawasaki (n=27), rashes (n=27)	Mechanical ventilation (n=14), antiviral (n=26), antibiotics (n=43), IVIG (n=15), systemic steroids (n=32), antimalarial (n=29)	Recovered (n=50) ICU admission (n=14)
Kainth, MK 2020	USA	HIC	65	Patients admitted in hospital	9.6	33	Travel history (NR) Contact history (n=33) Comorbidities (n=30)	Fever (n=57), cough (n=32), fatigue (n=26), dyspnea (n=34), GI symptoms (n=40), nasal symptoms or sore throat	Mechanical ventilation (n=5), antiviral (n=4), antibiotics (n=13), systemic steroids	Recovered (n=57) ICU admission (n=22) Death (n=1)

								(n=22), neurological (n=21), rashes (n=5)	(n=7), antimalarial (n=13), inotropes (n=8)	
Kaushik, S 2020	USA	HIC	33	Patients admitted in hospital	9.8	20	Travel history (NR) Contact history (n=13) Comorbidities (n=16)	Fever (n=31), dyspnea (n=11), diarrhea (n=16), nausea or vomiting (n=23), abdominal pain (n=21), neurological (n=4), Kawasaki signs (n=33), rashes (n=7), conjunctivitis (n=12)	Mechanical ventilation (n=5), antiviral (n=7), antibiotics (n=32), systemic steroids (n=17), IVIG (n=18), inotropes (n=17)	Recovered (n=32) ICU admission (n=33) Death (n=1)
Kest, H 2020	USA	HIC	3	Patients admitted in hospital	8.3	1	Travel history (n=0) Contact history (n=1) Comorbidities (n=0)	Fever (n=3), fatigue (n=1), GI symptoms (n=3), neurological (n=2), Kawasaki signs (n=3), rashes (n=3), conjunctivitis (n=3)	Mechanical ventilation (n=2), antibiotics (n=2), IVIG (n=2), systemic steroids (n=3), antimalarial (n=1), inotropes (n=3)	Recovered (n=3) ICU admission (n=3)
Korkmaz, MF 2020	Turkey	LMIC	81	Patients admitted in hospital	9.2	48	Travel history (NR) Contact history (n=75) Comorbidities (n=4)	Fever (n=47), cough (n=42), fatigue (n=15), dyspnea (n=2), GI symptoms (n=3), nasal symptoms or sore throat (n=7), neurological (n=11), asymptomatic (n=17)	Antibiotics (n=42), systemic steroids (n=3)	Recovered (n=81) ICU admission (n=2)
Meslin, P 2020	France	HIC	6	Patients admitted in hospital	0.1	3	Travel history (NR) Contact history (n=4) Comorbidities (n=2)	Fever (n=6), cough (n=1), fatigue (n=1), nausea or vomiting (n=1), nasal symptoms (n=2)	NR	Recovered (n=6) ICU admission (n=0)
Moraleda, C 2020	Spain	HIC	31	Patients admitted in hospital	7.8	18	Travel history (NR) Contact history (n=16)	Fever (n=30), cough (n=11), fatigue (n=5), dyspnea (n=8), GI symptoms (n=27), sore throat or nasal symptoms (n=8),	Mechanical ventilation (n=6), antiviral (n=9), antibiotics (n=28), systemic steroids (n=21), IVIG (n=20)	Recovered (n=29) ICU admission (n=20) Death (n=1)

							Comorbidities (n=10)	neurological (n=10), Kawasaki signs (n=13), rashes (n=23)		
Foong, Ng 2020	UK	HIC	3	Patients admitted in hospital	15.3	2	Travel history (n=0) Contact history (n=0) Comorbidities (n=1)	GI symptoms (n=3), Kawasaki signs (n=3), rashes (n=3), conjunctivitis (n=3)	Antibiotics (n=3), systemic steroids (n=1), IVIG (n=1), inotropes (n=2)	Recovered (n=3) ICU admission (n=3)
Pacquette, D 2020	Canada	HIC	938	National surveillance	NR	463	Travel history (n=106) Contact history (n=305) Comorbidities (NR)	NR	NR	NR
Parri, N 2020	Italy	HIC	130	Data collected through large collaborative networks	5.8	73	Travel history (n=0) Contact history (n=70) Comorbidities (N=34)	Fever (n=67), cough (n=54), dyspnea (n=17), diarrhea (n=10), sore throat or nasal symptoms (n=25), nausea or vomiting (n=15), neurological (n=3), asymptomatic (n=17)	Mechanical ventilation (n=2)	Recovered (n=130) ICU admission (n=9)
Perez, A 2020	USA	HIC	2	Patients admitted in hospital	16.5	1	Travel history (n=0) Contact history (n=0) Comorbidities (N=0)	Fever (n=1), fatigue (n=1), diarrhea (n=1), nausea or vomiting (n=1), abdominal pain (n=1)	NR	Recovered (n=2) ICU admission (n=0)
Sarangi, B 2020	India	LMIC	50	Patients presenting to in hospital	NR	28	Travel history (n=1) Contact history (n=45) Comorbidities (N=2)	Fever (n=17), cough (n=8), fatigue (n=4), diarrhea (n=2), sore throat or nasal symptoms (n=7), neurological (n=2), rashes (n=1),	NR	Recovered (n=50) ICU admission (n=0)

								asymptomatic (n=29), conjunctivitis (n=1)		
Sadiq, M 2020	Pakistan	LMIC	8	Patients admitted in hospital	9.5	7	Travel history (n=0) Contact history (n=8) Comorbidities (N=0)	Fever (n=8), fatigue (n=6), GI symptoms (n=6), neurological (n=3), Kawasaki signs (n=8), rashes (n=5), conjunctivitis (n=7)	Mechanical ventilation (n=1), systemic steroids (n=3), IVIG (n=6),	Recovered (n=5) ICU admission (n=2) Death (n=1)
Swann, OV 2020	UK	HIC	651	Patients admitted in hospitals	5.8	367	Travel history (NR) Contact history (NR) Comorbidities (n=276)	Fever (n=429), cough (n=231), fatigue (n=141), dyspnea (n=172), diarrhea (n=98), nausea or vomiting (n=179), abdominal pain (n=107), sore throat or nasal symptoms (n=132), neurological (n=148), kawasaki signs (n=52), rashes (n=87), conjunctivitis (n=26)	Mechanical ventilation (n=58), antiviral (n=38), antibiotics (n=415), systemic steroids (n=77), inotropes (n=47)	Recovered (n=557) ICU admission (n=116) Death (n=6)
Antunez-Montez, O 2020	5 Latin american Countries	LMIC	409	Patients presenting to hospitals	5.05	222	Travel history (n=0) Contact history (n=232) Comorbidities (N=83)	Fever (n=238), dyspnea (n=102), GI symptoms (n=101), sore throat or nasal symptoms (n=244), neurological (n=48), kawasaki signs (n=95), rashes (n=95), conjunctivitis (n=95), asymptomatic (n=49)	Mechanical ventilation (n=29), antiviral (n=14), IVIG (n=40), systemic steroids (n=63), antimalarial (n=10)	Recovered (n=392) ICU admission (n=52) Death (n=17)
Capone, C 2020	USA	HIC	33	Patients admitted in hospital	8.8	20	Travel history (NR) Contact history (NR) Comorbidities (N=7)	Fever (n=33), cough (n=17), dyspnea (n=17), diarrhea (n=32), nausea or vomiting (n=32), abdominal pain (n=32), neurological (n=19), kawasaki signs (n=21),	Mechanical ventilation (n=6), IVIG (n=33), systemic steroids (n=23), aspirin (n=29)	Recovered (n=33) ICU admission (n=52) Death (n=17)

								rashes (n=21), conjunctivitis (n=21)		
Cura Yayla, B 2020	Turkey	LMIC	220	Patients admitted in hospitals	9.3	106	Travel history (NR) Contact history (n=217) Comorbidities (N=21)	Fever (n=89), cough (n=79), fatigue (n=36), dyspnea (n=9), diarrhea (n=17), nausea or vomiting (n=9), sore throat or nasal symptoms (n=37), neurological (n=20), conjunctivitis (n=3), asymptomatic (n=55)	Mechanical ventilation (n=3), antiviral (n=6), antibiotics (n=40), antimalarial (n=1)	Recovered (n=209) ICU admission (n=3) Death (n=2)
Davies, P 2020	UK	HIC	78	Patients admitted to intensive care units	11.0	52	Travel history (NR) Contact history (NR) Comorbidities (N=17)	Fever (n=78), diarrhea (n=50), nausea or vomiting (n=49), abdominal pain (n=48), kawasaki signs (n=68), rashes (n=35), conjunctivitis (n=23)	Mechanical ventilation (n=36), antiviral (n=1), antibiotics (n=78), systemic steroids (n=57), IVIG (n=59), inotropes (n=65), aspirin (n=45)	Recovered (n=76) ICU admission (n=78) Death (n=2)
de Farias, E 2020	Brazil	LMIC	11	Patients admitted in hospital	5.4	9	Travel history (NR) Contact history (NR) Comorbidities (N=8)	Fever (n=11), kawasaki signs (n=5), rashes (n=5), conjunctivitis (n=5)	Mechanical ventilation (n=7), inotropes (n=7)	Recovered (n=9) ICU admission (n=11) Death (n=2)
Dima, M 2020	Romania	HIC	3	Patients admitted in hospital	Newborns	2	Travel history (n=0) Contact history (n=3) Comorbidities (N=1)	Fever (n=3), cough (n=1), fatigue (n=3), dyspnea (n=3)	antibiotics (n=1), IVIG (n=1), inotropes (n=47)	Recovered (n=3) ICU admission (n=0) Death (n=0)
Han, M 2020	Korea	HIC	91	Patients admitted in 20 hospitals	10.0	53	Travel history (n=15) Contact history (n=72)	Fever (n=27), cough (n=37), fatigue (n=12), dyspnea (n=5), diarrhea (n=11), nausea or vomiting (n=6),	antiviral (n=13), antimalarial (n=2)	Recovered (n=91) ICU admission (n=0) Death (n=0)

							Comorbidities (N=6)	abdominal pain (n=6), sore throat or nasal symptoms (n=8), neurological (n=12), asymptomatic (n=20)		
Ibrahim, L 2020	Australia	HIC	4	Patients presenting to hospital	13.1	1	Travel history (n=3) Contact history (n=4) Comorbidities (N=1)	Fever (n=1), cough (n=2), fatigue (n=4), diarrhea (n=1), sore throat or nasal symptoms (n=2), neurological (n=3)	Only supportive care	Recovered (n=4) ICU admission (n=0) Death (n=0)
Kilani, M 2020	Jordan	LMIC	61	Patients admitted in hospital	6.9		Travel history (n=0) Contact history (NR) Comorbidities (N=2)	Fever (n=11), cough (n=7), fatigue (n=12), diarrhea (n=10), sore throat or nasal symptoms (n=21), neurological (n=12), rashes (n=5), conjunctivitis (n=2), asymptomatic (n=27)	antimalarial (n=6)	Recovered (n=61) ICU admission (n=0) Death (n=0)
Okarska-Napierala, P 2020	Poland	HIC	9	Nationwide survey	10.35	4	Travel history (n=0) Contact history (n=2) Comorbidities (N=2)	Fever (n=9), cough (n=2), fatigue (n=2), dyspnea (n=2), diarrhea (n=5), nausea or vomiting (n=5), abdominal pain (n=5), neurological (n=5), kawasaki signs (n=9), rashes (n=6), conjunctivitis (n=4)	IVIG (n=7), systemic steroids (n=7), aspirin (n=6)	Recovered (n=4) ICU admission (n=1) Death (n=0)
Mahmoudi, S 2020	Iran	LMIC	35	Patients admitted in hospital	7.5	22	Travel history (n=0) Contact history (n=29) Comorbidities (N=18)	Fever (n=27), cough (n=28), fatigue (n=5), dyspnea (n=10), diarrhea (n=9), nausea or vomiting (n=10), abdominal pain (n=4), sore throat or nasal symptoms (n=3), neurological (n=7)	antiviral (n=30), antibiotics (n=35), antimalarial (n=22)	Recovered (n=31) ICU admission (n=7) Death (n=4)

Mamishi, S 2020	Iran	LMIC	45	Patients admitted in 3 hospitals	8.0	24	Travel history (n=0) Contact history (n=14) Comorbidities (N=6)	Fever (n=41), cough (n=16), fatigue (n=11), dyspnea (n=10), diarrhea (n=16), nausea or vomiting (n=23), abdominal pain (n=26), sore throat or nasal symptoms (n=7), kawasaki signs (n=31), rashes (n=24), conjunctivitis (n=23)	IVIg (n=18), systemic steroids (n=27),	Recovered (n=40) ICU admission (n=5) Death (n=5)
Mithal, L 2020	USA	HIC	18	Patients presenting to hospital	0.134	7	Travel history (n=0) Contact history (n=14) Comorbidities (N=0)	Fever (n=14), cough (n=8), fatigue (n=5), dyspnea (n=1), diarrhea (n=4), nausea or vomiting (n=4), sore throat or nasal symptoms (n=5), rashes (n=1), conjunctivitis (n=2), asymptomatic (n=1)	antibiotics (n=6)	Recovered (n=18) ICU admission (n=0) Death (n=0)
Newman, A 2020	USA	HIC	4	Patients admitted in hospital	10.9	4	Travel history (n=0) Contact history (n=0) Comorbidities (N=4)	Fever (n=3), cough (n=4), fatigue (n=2), dyspnea (n=1), nausea or vomiting (n=1), abdominal pain (n=1), sore throat or nasal symptoms (n=1)	Mechanical ventilation (n=1), antiviral (n=1), antibiotics (n=1), antimalarial (n=1)	Recovered (n=4) ICU admission (n=1) Death (n=0)
Torres, J 2020	Chile	HIC	27	Patients admitted in hospital	6.5	14	Travel history (n=0) Contact history (n=9) Comorbidities (N=7)	Fever (n=27), cough (n=7), diarrhea (n=17), nausea or vomiting (n=13), abdominal pain (n=17), kawasaki signs (n=18), rashes (n=14), conjunctivitis (n=13)	Mechanical ventilation (n=12), antibiotics (n=24), IVIG (n=19), systemic steroids (n=17) inotropes (n=12), aspirin (n=17)	Recovered (n=27) ICU admission (n=16) Death (n=0)
Saha, S 2020	Bangladesh	LMIC	26		0.02	16	Travel history (n=0)	Fever (n=14)		Recovered (n=12)

				Patients admitted in hospital			Contact history (n=8) Comorbidities (N=17)		Mechanical ventilation (n=25)	ICU admission (n=0) Death (n=4)
Gale, C 2020	UK	HIC	66	Population based surveillance study	Newborns	27	Travel history (n=0) Contact history (n=34) Comorbidities (N=18)	Fever (n=35), cough (n=11), fatigue (n=33), dyspnea (n=24), diarrhea (n=4), nausea or vomiting (n=33), sore throat or nasal symptoms (n=26), neurological (n=5), rashes (n=2), asymptomatic (n=7)	Mechanical ventilation (n=3), antiviral (n=2), antibiotics (n=46), IVIG (n=1), systemic steroids (n=2), inotropes (n=1)	Recovered (n=58) ICU admission (n=4) Death (n=1)
Schwartz, D 2020	Iran	LMIC	17	Patients admitted in hospital	Newborns	10	Travel history (n=0) Contact history (n=11) Comorbidities (NR)	Fever (n=4), cough (n=3), fatigue (n=17), dyspnea (n=12), diarrhea (n=1), abdominal pain (n=1), asymptomatic (n=2)	Mechanical ventilation (n=4), antibiotics (n=5), antimalarial (n=1), inotropes (n=1)	Recovered (n=7) ICU admission (n=4) Death (n=2)
Schwartz, S 2020	USA	HIC	4	Patients admitted in hospital	12.0	2	Travel history (n=0) Contact history (n=0) Comorbidities (N=3)	Fever (n=4), cough (n=3), fatigue (n=3), dyspnea (n=3), diarrhea (n=2), nausea or vomiting (n=3), abdominal pain (n=2), sore throat or nasal symptoms (n=2), rashes (n=1)	Mechanical ventilation (n=2), antiviral (n=2), systemic steroids (n=1), inotropes (n=1)	Recovered (n=4) ICU admission (n=4) Death (n=0)
Jain, S 2020	India	LMIC	23	Patients admitted in hospital	7.3	11	Travel history (n=0) Contact history (n=8) Comorbidities (NR)	Fever (n=23), cough (n=0), fatigue (n=3), dyspnea (n=11), diarrhea (n=15), nausea or vomiting (n=15), abdominal pain (n=12), sore throat or nasal symptoms (n=0), kawasaki signs (n=23),	Mechanical ventilation (n=9), IVIG (n=15), systemic steroids (n=22) inotropes (n=15)	Recovered (n=22) ICU admission (n=15) Death (n=1)



								rashes (n=14), conjunctivitis (n=11)		
Webb, K 2020	South Africa	LMIC	23	Patients admitted in hospital	6.6	6	Travel history (n=0) Contact history (n=3) Comorbidities (N=8)	Fever (n=23), cough (n=0), fatigue (n=8), dyspnea (n=10), diarrhea (n=13), nausea or vomiting (n=0), abdominal pain (n=15), sore throat or nasal symptoms (n=0), neurological (n=5), kawasaki signs (n=16), rashes (n=20), conjunctivitis (n=15)	Mechanical ventilation (n=6), IVIG (n=23), systemic steroids (n=15) inotropes (n=9), aspirin (n=17)	Recovered (n=15) ICU admission (n=12) Death (n=0)
van der Zalm, M 2020	South Africa	LMIC	159	Patients presenting to hospital	4.5	88	Travel history (NR) Contact history (NR) Comorbidities (N=33)	Fever (n=50), cough (n=72), fatigue (n=0), dyspnea (n=27), diarrhea (n=22), nausea or vomiting (n=18), abdominal pain (n=0), sore throat or nasal symptoms (n=18), neurological (n=20), kawasaki signs (n=3), rashes (n=6)	Mechanical ventilation (n=4)	Recovered (n=149) ICU admission (n=11) Death (n=1)
Sethuraman, U 2020	USA	HIC	34	Patients presenting to emergency department of the hospital	6.0	16	Travel history (n=0) Contact history (n=6) Comorbidities (N=11)	Fever (n=30), cough (n=0), fatigue (n=0), dyspnea (n=1), diarrhea (n=13), nausea or vomiting (n=14), abdominal pain (n=20), sore throat or nasal symptoms (n=9), rashes (n=19), conjunctivitis (n=9)	Mechanical ventilation (n=8), IVIG (n=24), inotropes (n=10), aspirin (n=24)	Recovered (n=34) ICU admission (n=24) Death (n=0)
Onal, P 2020	Turkey		37		10.0	20	Travel history (n=0)	Fever (n=26), cough (n=28), fatigue (n=18), dyspnea (n=13), diarrhea	Mechanical ventilation (n=1), antivirals (n=22)	Recovered (n=37)

		LMIC		Patients presenting to hospital			Contact history (n=29) Comorbidities (N=10)	(n=4), nausea or vomiting (n=8), abdominal pain (n=12), sore throat or nasal symptoms (n=12), neurological (n=11), asymptomatic (n=2)	antibiotics (n=24), IVIG (n=19), antimalarials (n=14)	ICU admission (n=10) Death (n=0)
Rabha, A 2020	Brazil	LMIC	115	Patients admitted in hospital	3.3	66	Travel history (n=0) Contact history (n=86) Comorbidities (N=30)	Fever (n=79), cough (n=59), fatigue (n=31), dyspnea (n=15), diarrhea (n=15), nausea or vomiting (n=20), abdominal pain (n=10), sore throat or nasal symptoms (n=17), rashes (n=5), neurological (n=23), asymptomatic (n=3)	antibiotics (n=24)	Recovered (n=115) ICU admission (n=12) Death (n=0)
Shahbaz nejad, L 2020	Iran	LMIC	10	Patients admitted in hospital	5.4	6	Travel history (n=0) Contact history (n=10) Comorbidities (N=0)	Fever (n=10), fatigue (n=2), dyspnea (n=2), diarrhea (n=7), nausea or vomiting (n=6), abdominal pain (n=9), sore throat or nasal symptoms (n=4), rashes (n=8), kawasaki signs (n=10), neurological (n=1), conjunctivitis (n=3)	Antivirals (n=1), antibiotics (n=10), IVIG (n=6), systemic steroids (n=2), antimalarial (n=8) inotropes (n=2), aspirin (n=3)	Recovered (n=9) ICU admission (n=3) Death (n=1)
Storch-de-Garcia, P 2020	Spain	HIC	39	Patients admitted in hospital	8.5	23	Travel history (n=0) Contact history (n=27) Comorbidities (N=5)	Fever (n=32), cough (n=20), fatigue (n=13), dyspnea (n=9), diarrhea (n=14), nausea or vomiting (n=15), abdominal pain (n=12), sore throat or nasal symptoms (n=18), rashes (n=10), kawasaki signs	Antivirals (n=13), antibiotics (n=25), IVIG (n=4), systemic steroids (n=15), antimalarial (n=15), aspirin (n=12)	Recovered (n=39) ICU admission (n=14) Death (n=0)

								(n=11), neurological (n=7), asymptomatic (n=7)		
García-Salido, A 2020	Spain	HIC	74	47 pediatric intensive care centres	7.7	46	Travel history (n=0) Contact history (n=13) Comorbidities (N=22)	Fever (n=61), cough (n=26), fatigue (n=48), dyspnea (n=23), diarrhea (n=33), nausea or vomiting (n=38), abdominal pain (n=33), sore throat or nasal symptoms (n=11), kawasaki signs (n=45), neurological (n=22)	Mechanical ventilation (n=18), antivirals (n=35), antibiotics (n=65), IVIG (n=25), systemic steroids (n=49), antimalarial (n=43), inotropes (n=37)	Recovered (n=65) ICU admission (n=74) Death (n=3)
Lima-setta, F 2020	Brazil	LMIC	56	17 pediatric intensive care centres	6.3	39	Travel history (n=0) Contact history (n=27) Comorbidities (N=11)	Fever (n=56), cough (n=16), fatigue (n=28), dyspnea (n=9), diarrhea (n=30), nausea or vomiting (n=21), abdominal pain (n=30), sore throat or nasal symptoms (n=17), rashes (n=38), kawasaki signs (n=26), neurological (n=27), conjunctivitis (n=26)	Mechanical ventilation (n=6), antivirals (n=5), antibiotics (n=33), IVIG (n=50), systemic steroids (n=31), aspirin (n=25)	Recovered (n=53) ICU admission (n=56) Death (n=1)
Zachariah, P 2020	USA	HIC	50	Patients admitted in hospital	11.5	27	Travel history (n=0) Contact history (n=9) Comorbidities (N=33)	Fever (n=40), cough (n=23), dyspnea (n=17), diarrhea (n=7), nausea or vomiting (n=7), abdominal pain (n=7), sore throat or nasal symptoms (n=6), neurological (n=3)	Mechanical ventilation (n=9), antivirals (n=4), IVIG (n=1), antimalarial (n=15)	Recovered (n=47) ICU admission (n=9) Death (n=1)
Kalamdani, P 2020	India	LMIC	12	Patients admitted in hospital	Newborns	8	Travel history (n=0) Contact history (n=12)	Fever (n=3), fatigue (n=2)	NR	Recovered (n=12) ICU admission (n=0) Death (n=0)

							Comorbidities (NR)			
Colomer, B 2020	Spain	HIC	40	Cases from a National Registry supported by the Spanish Society of Neonatology	Newborns	25	Travel history (n=0) Contact history (n=36) Comorbidities (N=15)	Fever (n=18), cough (n=9), fatigue (n=4), dyspnea (n=7), diarrhea (n=2), nausea or vomiting (n=4), sore throat or nasal symptoms (n=5), rashes (n=1), asymptomatic (n=10)	Antivirals (n=1), antibiotics (n=8), IVIG (n=19), antimalarial (n=5)	Recovered (n=40) ICU admission (n=8) Death (n=0)
Dhanalakshmi, K 2020	India	LMIC	19	Patients admitted in hospital	7.2	8	Travel history (n=0) Contact history (n=0) Comorbidities (N=1)	Fever (n=19), dyspnea (n=8), diarrhea (n=3), nausea or vomiting (n=6), abdominal pain (n=8), sore throat or nasal symptoms (n=9), rashes (n=12), neurological (n=6), kawasaki signs (n=7), conjunctivitis (n=9)	Antibiotics (n=19), IVIG (n=15), systemic steroids (n=11) inotropes (n=6), aspirin (n=16)	Recovered (n=19) ICU admission (n=12) Death (n=0)
Falah, N 2020	Pakistan	LMIC	10	Patients admitted in 2 hospitals	6.0	8	Travel history (n=0) Contact history (n=2) Comorbidities (N=0)	Fever (n=10), cough (n=2), fatigue (n=4), dyspnea (n=2), diarrhea (n=3), abdominal pain (n=5), sore throat or nasal symptoms (n=6), rashes (n=8), neurological (n=6), kawasaki signs (n=5), conjunctivitis (n=9)	Antivirals (n=1), antibiotics (n=6), IVIG (n=9), systemic steroids (n=3) inotropes (n=4), aspirin (n=7)	Recovered (n=10) ICU admission (n=7) Death (n=0)
Zheng, G 2020	China	LMIC	52	Patients admitted in 11 hospitals	8.5	28	Travel history (n=52) Contact history (n=42) Comorbidities (N=0)	Fever (n=21), cough (n=25), fatigue (n=5), diarrhea (n=1), nausea or vomiting (n=1), abdominal pain (n=1), sore throat or nasal	Antivirals (n=43), antibiotics (n=25)	Recovered (n=52) ICU admission (n=0) Death (n=0)

								symptoms (n=3), asymptomatic (n=13)		
Almoosa, Z 2020	Saudi Arabia	HIC	10	Patients admitted in 2 hospitals	7.6	5	Travel history (n=0) Contact history (n=6) Comorbidities (N=5)	Fever (n=10), fatigue (n=5), dyspnea (n=3), diarrhea (n=7), nausea or vomiting (n=9), abdominal pain (n=9), Kawasaki signs (n=10), rashes (n=10), conjunctivitis (n=9)	Mechanical ventilation (n=2), antivirals (n=7), antibiotics (n=10), IVIG (n=10), antimalarials (n=10), systemic steroids (n=17) inotropes (n=5), aspirin (n=4)	Recovered (n=8) ICU admission (n=8) Death (n=2)
Alsharrah, D 2020	Kuwait	HIC	134	Patients admitted in hospital	8.7	74	Travel history (n=19) Contact history (n=110) Comorbidities (N=15)	Fever (n=75), cough (n=100), asymptomatic (n=91)	Antibiotics (n=11)	Recovered (n=133) ICU admission (n=0) Death (n=0)
Prata-Barbosa, A 2020	Brazil	LMIC	79	19 pediatric intensive care centres	4.8	43	Travel history (NR) Contact history (NR) Comorbidities (N=32)	Fever (n=59), cough (n=40), fatigue (n=35), dyspnea (n=23), diarrhea (n=16), nausea or vomiting (n=16), abdominal pain (n=16), sore throat or nasal symptoms (n=17), kawasaki signs (n=5)	Mechanical ventilation (n=14), antivirals (n=34), antibiotics (n=60), systemic steroids (n=18) inotropes (n=12), aspirin (n=17)	Recovered (n=71) ICU admission (n=79) Death (n=2)
Bayesheva, D 2020	Kazakhstan	LMIC	650	Data was reported from the official medical electronic database of the Country	7.1	366	Travel history (NR) Contact history (NR) Comorbidities (N=76)	Fever (n=59), cough (n=96), fatigue (n=7), dyspnea (n=12), diarrhea (n=13), asymptomatic (n=315), sore throat or nasal symptoms (n=119), rashes (n=1)	Mechanical ventilation (n=3)	Recovered (n=650) ICU admission (n=6) Death (n=0)
Bustos-Cordova, E 2020	Mexico		50		5.9	35	Travel history (NR)	Fever (n=35), cough (n=22), fatigue (n=21), dyspnea (n=10), diarrhea		Recovered (n=50)

		LMIC		Patients presenting to tertiary care centre outpatient department			Contact history (NR) Comorbidities (N=26)	(n=11), nausea or vomiting (n=11), abdominal pain (n=11), sore throat or nasal symptoms (n=25), rashes (n=4), kawasaki signs (n=1), conjunctivitis (n=11)	Mechanical ventilation (n=2), antibiotics (n=15)	ICU admission (n=2) Death (n=1)
Bhavsar, M 2020	USA	HIC	81	Patients admitted in 12 hospitals in the city	14.7	40	Travel history (NR) Contact history (NR) Comorbidities (N=21)	Fever (n=63), cough (n=38), diarrhea (n=26), nausea or vomiting (n=26), abdominal pain (n=26), neurological (n=7), rashes (n=8), conjunctivitis (n=7)	Mechanical ventilation (n=3), antivirals (n=6), IVIG (n=1), systemic steroids (n=15), antimalarials (n=10), inotropes (n=9), aspirin (n=1)	Recovered (n=71) ICU admission (n=28) Death (n=1)
Montoya, J 2020	Peru	LMIC	69	Cancer patients admitted to tertiary care centres	6.0	44	Travel history (n=0) Contact history (n=20) Comorbidities (N=69)	Fever (n=12), cough (n=12), dyspnea (n=11), diarrhea (n=3), nausea or vomiting (n=3), abdominal pain (n=2), sore throat or nasal symptoms (n=11), rashes (n=1), asymptomatic (n=37)	Antibiotics (n=9), interferon (n=9), systemic steroids (n=9)	Recovered (n=56) ICU admission (n=3) Death (n=7)

NR= Not reported

LMIC: Low and middle-income Country, HIC: High-income Country

**Table 4: Age Distribution from 91 studies of disease prevalence**

<b>Age</b>	<b>Frequency (%)</b>
0 to ≤5 years	3149 (36.9)
>5 to ≤10 years	1668 (19.6)
>10 years	3712 (43.5)
Total	8529

**Table 5: Age Distribution from 67 case series of clinical features and outcomes**

	Age bands		
	0 to ≤5 years	>5 to ≤10 years	>10 years
<b>Clinical features</b> Frequency (%)			
Fever	99/181 (54.7)	87/133 (65.4)	60/132 (45.4)
Asymptomatic	43/1053 (4.0)	45/654 (7.0)	48/880 (5.4)
Cough	46/181 (25.4)	33/133 (24.8)	23/132 (17.4)
Nasal symptoms/sore throat	22/181 (12.1)	16/133 (12.0)	5/132 (3.8)
Dyspnea	15/181 (8.3)	10/133 (7.5)	13/132 (9.9)
Diarrhea	29/181 (16.0)	48/133 (36.1)	36/132 (27.3)
Nausea and vomiting	29/181 (16.0)	40/133 (30.0)	33/132 (25.0)
Abdominal pain	27/181 (15.0)	47/133 (35.3)	30/132 (22.7)
Fatigue	13/181 (7.2)	7/133 (5.2)	5/132 (3.8)
Neurological signs	8/181 (4.4)	21/133 (15.8)	16/132 (12.1)
Kawasaki signs	15/181 (8.3)	24/133 (18.0)	8/132 (6.0)
Rash	24/181 (13.2)	27/133 (20.3)	20/132 (15.1)
Conjunctivitis	23/181 (12.7)	8/133 (6.0)	18/132 (13.6)
<b>Outcomes</b> Frequency (%)			
Recovered	176/196 (89.8)	119/127 (93.7)	146/160 (91.2)
Severe/ICU admission	188/1708 (11.0)	102/834 (12.2)	131/1134 (11.6)
Not yet discharged	19/196 (9.7)	6/127 (4.7)	8/160 (5.0)
Deaths	1/1360 (0.07)	2/742 (0.3)	11/908 (1.2)



**Table 6: Clinical symptoms among MISC from 31 studies**

<b>Characteristics</b>	<b>Events/total patients</b>	<b>Mean proportion % (95% CI)</b>
Comorbidity	264/1182	24.7 (20.1-29.9)
Fever	1008/1208	94.2 (88.7-97.1)
Kawasaki shock/sign	708/1208	68.0 (56.0-78.1)
Abdominal pain	313/548	58.1 (49.6-66.1)
Diarrhea	308/563	54.0 (45.9-61.9)
Conjunctivitis	461/1163	49.2 (40.2-58.2)
Nausea/vomiting	280/563	48.7 (38.5-59.0)
Rashes	540/1163	46.5 (43.4-49.7)
Dyspnea	220/1181	22.4 (17.1-28.8)
Neurological symptoms	184/1206	19.0 (12.8-27.3)
Fatigue	104/626	16.3 (9.6-26.3)
Cough	94/766	15.1 (10.1-22.2)
Nasal symptoms	310/1124	12.1 (6.9-20.5)
Pharyngeal erythema	298/1099	10.3 (5.5-18.6)
<b>Clinical management</b>		
IVIG	618/1207	69.1 (55.1-80.4)
Inotropes	255/934	45.5 (34.3-57.2)
Antibiotics	317/1021	38.6 (19.9-61.3)
Aspirin	201/951	31.7 (19.5-47.1)
Systemic steroids	522/1207	52.9 (41.6-63.8)
Mechanical ventilation	208/1204	20.2 (13.9-28.4)
Antimalarial	60/954	7.3 (3.0-16.8)
Antiviral	63/1015	6.9 (3.8-12.4)
<b>Clinical outcomes</b>		
Recovered	1099/1208	91.8 (86.8-95.1)
ICU admission	638/1208	47.1 (43.1-51.1)
Deaths	39/1208	4.8 (3.6-6.3)

**Table 7: Clinical symptoms among neonates from 11 studies**

<b>Characteristics</b>	<b>Events/total patients</b>	<b>Mean proportion % (95% CI)</b>
Comorbidity	53/184	28.0 (16.5-43.3)
Fever	90/184	47.6 (35.8-59.7)
Dyspnea	50/146	39.1 (22.1-59.2)
Inability to feed/Lethargy	62/158	35.4 (16.7-59.9)
Nausea/vomiting	38/146	34.4 (25.9-44.0)
Nasal symptoms	36/146	29.9 (22.3-38.9)
Cough	28/158	19.2 (13.6-26.3)
Asymptomatic	21/146	16.6 (11.2-23.9)
Pharyngeal erythema	8/146	11.4 (4.8-24.9)
Diarrhea	8/146	7.6 (4.1-13.4)
Neurological symptoms	5/146	7.3 (3.8-13.5)
Rashes	3/146	5.3 (2.4-11.0)
<b>Clinical management</b>		
Antibiotics	86/166	48.7 (39.6-58.0)
Mechanical ventilation	8/140	10.7 (5.2-20.8)
Antimalarial	6/140	9.6 (5.1-17.6)
Antiviral	3/140	6.4 (3.0-12.9)
IVIG	2/140	6.3 (2.4-15.4)
Inotropes	2/140	6.3 (2.4-15.4)
Systemic steroids	2/140	6.3 (2.4-15.4)
<b>Clinical outcomes</b>		
Recovered	146/184	77.8 (55.7-90.7)
ICU admission	22/184	16.0 (7.6-30.7)
Deaths	7/184	8.8 (4.9-15.3)

**Table 8: Quality assessment of included studies (N=129)**

Study and year	Was the study question or objective clearly stated?	Was the study population clearly and fully described, including a case definition?	Were the cases consecutive?	Were the subjects comparable?	Was the intervention clearly described?	Were the outcome measures clearly defined, valid, reliable, and implemented consistently across all study participants?	Was the length of follow-up adequate?	Were the statistical methods well described?	Were the results well described?	Overall Quality (Good, fair or poor)
Qiu, H 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Wang, D 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Zheng, F 2020	Yes	Yes	No	Yes	NA	Yes	No	Yes	No	5
Xia, W 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	No	6
Feng, K 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	No	6
Cai, J 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	No	6
Wei, M 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Su, L 2020	Yes	Yes	No	Yes	NA	Yes	No	Yes	Yes	6
Zhou, Y 2020	Yes	Yes	No	Yes	NA	Yes	No	Yes	Yes	6
Sun, D 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Liu, W 2020	Yes	Yes	No	Yes	NA	Yes	No	Yes	Yes	6
Hu, Z 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Liu, M 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Liu, H 2020	Yes	No	Yes	Yes	NA	Yes	No	Yes	No	5
Lou, MX 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Zhang, T 2020	Yes	No	No	Yes	NA	Yes	Yes	Yes	Yes	6
Li, Y 2020	Yes	Yes	No	Yes	NA	Yes	No	Yes	Yes	6
Ji, LN 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Tagarro, A 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Chen, A 2020	Yes	Yes	No	Yes	NA	Yes	No	Yes	No	5
See, KC 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8

Zhang, GX 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Cheng, X 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Tang, A 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Zhu, L 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Tan, X 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Ma, H 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	7
Shen, Q 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	7
Schwierzeck, V 2020	Yes	No	Yes	Yes	NA	Yes	No	Yes	No	5
Jiang, JY 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Riphagen, S 2020	Yes	No	Yes	Yes	NA	Yes	Yes	Yes	Yes	7
Shekerdemian, LS 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Verdoni, L 2020	Yes	No	Yes	Yes	NA	Yes	Yes	Yes	Yes	7
Rahimzadeh, G 2020	Yes	No	Yes	Yes	NA	Yes	Yes	Yes	Yes	7
Tullie, L 2020	Yes	No	No	Yes	NA	Yes	Yes	Yes	Yes	6
Sajid, MI 2020	Yes	Yes	Yes	Yes	NA	Yes	No	Yes	Yes	7
Zhang, ZJ 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	7
Zeng, L 2020	Yes	Yes	Yes	Yes	NA	Yes	No	Yes	Yes	7
Salvatori, G 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Zhang, X 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Wu, HP 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Zhang, H 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Chao, JY 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Eghbali, A 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Chiotos, K 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Flores, V 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
García-Salido, A 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Cabrero-Hernández, M 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Waltuch, T 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7

Kakuya, F 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Soltani, J 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Toubiana, J 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Foster, CE 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
White, A 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Lin, EE 2020	Yes	Yes	No	Yes	NA	Yes	No	Yes	No	5
Gao, Y 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Harman, K 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Blondiaux, E 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Pouletty, M 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Whittaker, E 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Cheung, EW 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Lu, X 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	7
CDC, COVID 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Gotzinger, F 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	7
Dong, Y 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	7
Abdel-Mannan, O 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Guha, A 2020	Yes	Yes	Yes	Yes	NA	Yes	No	Yes	Yes	7
Bhumbra, S 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Bisogno, G 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Blumfield, E 2020	Yes	Yes	No	Yes	NA	Yes	No	Yes	Yes	6
Derespina, KR 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Feldstein, LR 2020	Yes	Yes	No	Yes	NA	Yes	No	Yes	Yes	6


George, S 2020	Yes	Yes	Yes	Yes	NA	Yes	No	Yes	Yes	7
Gonzalez-Cortes, R 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Kainth, MK 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Kaushik, S 2020	Yes	Yes	Yes	Yes	NA	Yes	No	Yes	Yes	7
Kest, H 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Korkmaz, MF 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Meslin, P 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Moraleda, C 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	No	6
Foong, Ng 2020	Yes	Yes	No	Yes	NA	Yes	No	Yes	Yes	6
Pacquette, D 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Parri, N 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	7
Perez, A 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	No	6
Sarang, B 2020	Yes	Yes	Yes	Yes	NA	Yes	No	Yes	No	6
Sadiq, M 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Swann, OV 2020	Yes	Yes	Yes	Yes	NA	Yes	No	Yes	Yes	7
Antunez-Montez, O 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Capone, C 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Cura Yayla, B 2020	Yes	Yes	Yes	Yes	NA	Yes	No	Yes	No	6
Davies, P 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
de Farias, E 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Dima, M 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Han, M 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Ibrahim, L 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Kilani, M 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Okarska-Napierala, P 2020	Yes	Yes	Yes	Yes	NA	Yes	No	Yes	No	6

Mahmoudi, S 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Mamishi, S 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Mithal, L 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Newman, A 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Torres, J 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Saha, S 2020	Yes	Yes	No	Yes	NA	Yes	No	Yes	No	5
Gale, C 2020	Yes	Yes	Yes	Yes	NA	Yes	No	Yes	No	6
Schwartz, D 2020	Yes	Yes	No	Yes	NA	Yes	No	Yes	No	5
Schwartz, S 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Jain, S 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Webb, K 2020	Yes	No	No	Yes	NA	Yes	No	Yes	Yes	5
van der Zalm, M 2020	Yes	Yes	Yes	Yes	NA	Yes	No	Yes	No	6
Sethuraman, U 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Onal, P 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Rabha, A 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8
Shahbaznejad, L 2020	Yes	No	No	Yes	NA	Yes	Yes	Yes	Yes	6
Storch-de-Garcia, P 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
García-Salido, A 2020	Yes	Yes	No	Yes	NA	Yes	No	Yes	Yes	6
Lima-setta, F 2020	Yes	Yes	No	Yes	NA	Yes	No	Yes	Yes	6
Zachariah, P 2020	Yes	Yes	No	Yes	NA	Yes	No	Yes	Yes	6
Kalamdani, P 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	7
Colomer, B 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	Yes	8

Dhanalakhmsi, K 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Falah, N 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Zheng, G 2020	Yes	No	Yes	Yes	NA	Yes	Yes	Yes	Yes	7
Almoosa, Z 2020	Yes	Yes	No	Yes	NA	Yes	Yes	Yes	Yes	7
Alsharrah, D 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	7
Prata-Barbosa, A 2020	Yes	Yes	Yes	Yes	NA	Yes	No	Yes	Yes	7
Bayesheva, D 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	7
Bustos-Cordova, E 2020	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	7
Bhavsar, M 2020	Yes	Yes	No	Yes	NA	Yes	No	Yes	Yes	6
Montoya, J 2020	Yes	No	Yes	Yes	NA	Yes	No	Yes	Yes	6



Figure1: Flow diagram of the systematic review process



**Figure 2: Global map of distribution of the studies included (N=129) in the review**



Created by Datawrapper <https://www.datawrapper.de/>